

ASAIM/led-Sci News

Newsletter of The American Society of Addiction Medicine

Inside

MED-SCI COVERAGE:

Quick Guide to the Med-Sci Program / 10

Map of the Hotel $\,$ / ${\bf S26}$

Exhibitors & Locations / **\$25**

Registration Desk Hours / **9**

Med-Sci Overview / 8

Ruth Fox Course for Physicians / 9

Buprenorphine Training Course / 24

NIAAA Symposium / **\$25**

NIDA Symposium / **\$25**

NIDA Symposium / 323

CSAT Symposium / **S26**

ASAM Awards / 5

State Meetings / **S26**

Ruth Fox Donor Reception / 22

Med-Sci Funders / **S26**

Visiting San Diego / 8

ASAM Med-Sci News is made possible by an unrestricted educational grant from Forest Pharmaceuticals, Inc.

www.asam.org

A Message from the President:

Welcome to ASAM's 37th Annual Conference

Elizabeth F. Howell, M.D., FASAM ASAM President



A s we gather again for ASAM's annual Medical-Scientific Conference, there is value in revisiting the path we have traveled to arrive at ASAM's current position as a Society.

Today, ASAM is a vibrant national organization of more than 3,000 members who are

dedicated to the belief that the treatment of addiction should be granted parity with the treatment of any other chronic medical disorder, that all physicians should receive education in addiction medicine, and that physicians who wish should be trained and board-certified in addiction medicine.

By contrast, in 1954, when our colleagues organized as the New York Medical Society on Alcoholism and held their first scientific meeting, fewer than 25 persons registered. At this year's ASAM Medical-Scientific Conference, we expect 1,000 registrants.

In 1954, there was no textbook of addiction medicine. In 2003, ASAM published the Third Edition of its *Principles of Addiction Medicine*, which in 1,600 pages reflects the vast body of scientific knowledge in this field.

In 1954, there were no national guidelines for determining the need for treatment and the appropriate level of care. Today, the *ASAM Patient Placement Criteria* are required or recommended by many states and by an increasing number of managed care organizations.

In 1954, there were no practice guidelines. Today, ASAM has published three guidelines — one in the prestigious *Journal of the American Medical Association*.

None of these accomplishments came easily. Each required the concerted effort of many dedicated individuals. But the results of their struggles have improved the lives of countless patients and their families. So too will our efforts to overcome the challenges that confront us today.

As we begin our 37th Annual Meeting, we salute you, the ASAM member, and invite you to continue to contribute ideas, suggestions, and efforts to your Society. For it is the responsibility of members to guide ASAM, to monitor its progress, to constructively criticize ASAM's efforts, and to constantly demand more of their Society, so that ASAM can achieve its full potential.

A Message from the President-Elect:

For ASAM Members: A Call to Action

Michael M. Miller, M.D., FASAM ASAM President-Elect



Welcome to San Diego! Thanks to all the presenters and staff, as well as to our host, the California Society of Addiction Medicine. Congratulations to CSAM for all it does to improve the practice of — and the practice environment for — addiction medicine.

As I begin the second year of my six-year term as your President-Elect, President and Immediate Past-President, I'd like to issue a Call to Action to you and to other ASAM members to get involved in ASAM and your local chapter. The challenge to ASAM's officers, Board, and Chapter Presidents is to identify specific channels for your involvement — especially channels to involve our newer and younger members. We must make better use of their energies, talents and passions to move ASAM's mission forward. Our members must never be in the position of saying, "I'd like to do something, but I don't know what or how."

One of the concerns we all share is the quality of medical care. Every ASAM physician is concerned about the plight of the person with addictive disease, worries about how misunderstood and stigmatized their illness is, and laments the barriers to care faced by most of our patients or potential patients. One way to close gaps in quality and access is to become involved in the public policy arena. Through the leadership of Kevin Kunz, M.D., and ASAM consultant James F. Callahan, D.P.A., we are working toward the goal of having an active public policy committee in every State chapter to coalesce the energies of ASAM members in each State, and to advocate with State and Federal agencies and legislators on behalf of our patients and our practices.

If your state chapter hasn't yet realized this vision, perhaps you are the person who'll step up to the plate and take on the challenge. Alexis Geier-Horan in ASAM's national office; ASAM Public Policy Committee co-chairs Mark Kraus, M.D., and Petros Levounis, M.D., as well as Kevin and Jim and Legislative Advocacy Committee chair Don Kurth, M.D., and all the ASAM officers are ready to assist any ASAM member who decides to direct his or her volunteer efforts into public policy.

(continued on page S2)

A Message from the Executive Vice President:

Strengthening ASAM by Meeting Members' Needs

What role should ASAM occupy in addiction modicing tion medicine today, and in the future? Those of you who are veteran members of ASAM may remember why you originally joined the Society, how you expected to benefit from membership, and how you expected ASAM to influence addiction medicine. No doubt some of your expectations have been realized and some have not.

When I first joined ASAM, I asked the Society's leaders and members about their "vision" for ASAM's future. Their responses were thought-provoking, and remain relevant today.

For example, several leaders described ASAM's principal challenge as engaging primary care medicine in addiction care and encouraging physicians to deal with addiction in their practices. They saw ASAM's role as educating physicians in primary care and in other specialties about the aspects of addiction that are relevant to their practices. In their view, ASAM should be a resource to the larger medical community.

Eileen McGrath, J.D.



ASAM's role in advocacy is a central part of the vision of some ASAM members I spoke with, who see ASAM as the natural leader of ongoing efforts to achieve parity and reduce stigma. They want ASAM to be at the forefront not only on "pocketbook" issues - which sometimes promote a self-serving appearance — but also to advocate for patients' unmet needs. These members point out that achieving parity will require bipartisan support at both the federal and state levels.

Some ASAM members proposed expanding ASAM's membership to include allied health professionals as a way to build unity in the field. Others were equally adamant that for addiction medicine to achieve ABMS recognition as a medical specialty, ASAM must remain focused on the needs of physicians in addiction medicine.

Listening to members' views and opinions is a vital part of my role at ASAM. My vision is that ASAM will succeed in advancing addiction medicine only by meeting the needs of its members.

As we begin another annual meeting, we have an opportunity once again to share our visions and our commitment to the Society. I look forward to speaking with many of you during Med-Sci. I also invite you to contact me at any time, by e-mail at EMCGRATH@ ASAM.ORG or by mail or phone at the address on the ASAM News masthead. You have my commitment to attend to your comments and suggestions. What is your vision for ASAM's future? How can ASAM best meet your needs?

A CALL TO ACTION (continued from page S1)

Another way to improve the quality of care is to get involved in other quality efforts in medicine, so as to bring the perspective of addiction medicine to the QI activities already underway. Therefore, the other part of my Call to Action is to invite you to participate in ASAM's Quality Improvement Council. Just in time for our 2006 Medical-Scientific Conference, we have opened a new QI page on the ASAM website (WWW.ASAM.ORG). I hope I can excite your passion for quality by highlighting several points that are discussed in greater length on that new web link:

- 1. The Institute of Medicine's 2005 report, Improving the Quality of Health Care for Mental and Substance-Use Conditions, is a follow-up to their highly publicized reports of 1999 and 2001 (Crossing the Quality Chasm). ASAM provided formal commentary on the report and contributed to the development of a proposed legislative agenda in response to the report. Both documents are posted at WWW.ASAM.ORG.
- 2. The AMA's Physician Consortium for Performance Measurement provides a conduit for ASAM to be involved in the development of performance measures that relate to the care of patients with substance use and chronic pain disorders.
- 3. ASAM is involved with the Joint Commission on Accreditation of Healthcare Organizations.
- 4. ASAM is working with the National Association of Addiction Treatment Providers (NAATP) to identify performance measures for addiction treatment organizations.
- 5. ASAM's ongoing work in developing Practice Guidelines has built

- sound scientific underpinnings to many areas of addiction practice.
- 6. ASAM's continuous development and expansion of the ASAM Patient Placement Criteria provides the field with an independently developed, non-commercial system for matching patients' needs to available services.
- 7. ASAM has connections to the American College of Medical Quality — a national medical specialty society of physicians who work full-time in the quality arena.

There's not much debate over the fact that the quality of addiction care in America needs improvement. I hope you agree with me that there is no group of physicians better positioned to address these quality improvement challenges than the members of the American Society of Addiction Medicine. After all, ASAM is no more than you and me — our individual efforts focused collectively toward achieving specific goals. If individual ASAM members like you don't participate in defining what constitutes quality care, what constitutes the optimum measurement systems for defining quality in medical practice, and for influencing what major organizations like JCAHO are doing in the quality arena, then we can be assured that some other entities — be they physician organizations or agencies of government or other non-physician groups — will do this work for us and impose their expectations upon us.

Please contact ASAM's Executive Vice President, Eileen McGrath, J.D., any ASAM officer, or the co-chairs of ASAM's Quality Improvement council (Dr. David Gastfriend and Dr. David Mee-Lee) to let us know that you are ready to "answer the call" and to contribute to ASAM's activities in the quality improvement arena.



March-April 2006 Volume 21, Number 2

Newsletter of The American Society of Addiction Medicine

Inside

ASAM at Work for You:

Report from the EVP / 2

From the President / 4

Med-Sci Program Guide / **7**

State Society/ Chapter News / **15**

Membership News / 21

Ruth Fox Fund / 22

Calendar / 24

Other News:

Addiction
Medicine News / 3

Policy Matters / 5

Clinical Briefing / 6

Notes from the Field / 12

See pages 7-11 for your handy guide to ASAM's Med-Sci Conference

www.asam.org



San Diego to Host ASAM's 37th Annual Med-Sci Conference

S an Diego, California, is the site of ASAM's 37th Annual Medical-Scientific Conference, to be held May 4-7th at the Sheraton Hotel and Marina. The conference welcomes ASAM members as well as non-member physicians, nurses, psychologists, counselors, students and residents. It features three full days of clinical and scientific programming, as well as ASAM's annual Business Meeting at 7:00 a.m. Friday, May 5th.

The Med-Sci Conference is preceded by the Ruth Fox Course for Physicians on Thursday, May 4th, also at the San Diego Sheraton Hotel and Marina. The educational activities conclude on Sunday, May 7th, with a Buprenorphine Training Course.

For additional information or to register, visit the ASAM website at WWW.ASAM.ORG or contact ASAM's Department of Meetings and Conferences at 301/656-3920. (Conference coverage continues on page 7)

New Educational Program Debuts at Med-Sci

A new ASAM medical education program on "Pharmacologic Therapies for Alcohol Dependence" will be offered for the first time during ASAM's Med-Sci Conference. Scheduled for Saturday evening, May 6th, the program will begin with dinner at 6:00 p.m., followed by the presentations from 7:00 to 9:00 p.m. Thanks to an unrestricted educational grant from Cephalon, Inc., there is no registration fee, but advance registration is required.

The program focuses on how to use ASAM's Patient Placement Criteria to successfully integrate new pharmacologic therapies with other components of addiction care. Faculty include noted addiction researcher and educator Charles P. O'Brien, M.D., Ph.D., of the University of Pennsylvania, a

recipient of ASAM's Distinguished Scientist Award. Dr. O'Brien will be joined by ASAM Criteria developers David Mee-Lee, M.D., and Gerald Shulman, M.A., FACATA, as well as other distinguished experts.

Chairman Marc Fishman, M.D., says the program — which is approved for 2 hours of Category 1 CME credit — is designed to be highly interactive and will include case vignettes and ample opportunity for audience-faculty interaction. In addition, each registrant will receive a program book that includes copies of the speakers' slides and other relevant material. For more information or to register, visit the ASAM website at www.asam.org or contact Angela Warner at awarner@asam.org.

REPORT FROM THE EVP

ASAM Certification Is a Valuable Credential

Eileen McGrath, J.D., Executive Vice President/CEO

s the recruitment advertise-and other publications testify, ASAM certification in addiction medicine has become a valuable credential. Certification signifies that an individual has mastered the scientific and clinical aspects of this emerging specialty, as demonstrated on a rigorous examination prepared by the National Board of Medical Examiners. Since the



Eileen McGrath, J.D.

exam first was offered in 1986, more than 4,000 physicians — including many of the nation's top addiction treatment professionals — have been certified by ASAM.

Physicians who wish to register for ASAM's 2006 Certification/Recertification Examination must complete and submit an application no later than April 30th. After the applications are reviewed, candidates will notified by mail as to whether they qualify to sit for the examination, which will be given on Saturday, December 9th, 2006, at three sites: Los Angeles, CA; New York, NY; and Atlanta, GA.

The fee for the examination is \$1,350 for ASAM members and \$1,600 for non-members. For more information, contact **ASAM Credentialing Director** Christopher M. Weirs, M.P.A., by email at CWEIR@ASAM.ORG or by phone at 301/656-3920, or visit the ASAM website at

WWW.ASAM.ORG. The website contains a downloadable application and information about qualifications to sit for the examination.

Also, watch for the May-June ASAM News for details about the Review Course in Addiction Medicine, to be offered Thursday, October 26th through Saturday, October 28th, at Chicago's Westin O'Hare Hotel. For more information or to register for the course, visit the ASAM website at www.ASAM.ORG.

See the May-June issue of ASAM News for details about ASAM's Review Course in Addiction Medicine, to be offered Thursday, October 26th through Saturday, October 28th, at the Westin O'Hare Hotel in Chicago.



JACKSONVILLE, FLORIDA — Chemical Dependence Facility seeking Board Certified Internal Medicine or Family Practice physician to join our team. ASAM Certification preferred. Competitive salary and excellent benefits, including 401K match. Email: mprovines@lakeviewhealth.com, or fax 954/491-4193.



American Society of **Addiction Medicine**

4601 North Park Ave., Suite 101 Chevy Chase, MD 20815

ASAM is a specialty society of physicians concerned about alcoholism and other addictions and who care for persons affected by those illnesses.

Officers

President

Elizabeth F. Howell, M.D., FASAM

Immediate Past President Lawrence S. Brown, Jr., M.D., M.P.H., FASAM

President-Elect

Michael M. Miller, M.D., FASAM, FAPA

Secretary

A. Kenison Roy III, M.D., FASAM

Treasurer

Donald J. Kurth, M.D., FASAM

Executive Vice President/CEO Eileen McGrath, J.D.

ASAM News

is an official publication of the American Society of Addiction Medicine. It is published six times a year. Please direct all inquiries to the Editor at ASAMNEWSLETTER@AOL.COM or phone 410/770-4866.

Chair, Publications Council Elizabeth F. Howell, M.D., FASAM

Newsletter Review Board LeClair Bissell, M.D. Sheila B. Blume, M.D., FASAM Max A. Schneider, M.D., FASAM

Founding Editor, 1985-1995 Lucy Barry Robe

Editor

Bonnie B. Wilford

Subscriptions

Free to ASAM members; \$99 a year (six issues) to nonmembers. To order, phone 1-800/844-8948 or fax 301/206-9789.

Advertising

Advertising rates and schedules are available on request. Please direct inquiries to the Editor at 410/770-4866 or email ASAMNEWSLETTER@AOL.COM.

Web Site

For members visiting ASAM's web site (WWW.ASAM.ORG), entrance to the on-line Membership Directory requires the Username "asam" and the password "asam" (in lower case letters).

DOD Study Finds MH, Substance Abuse **Problems Among Returning Iraq Vets**

Thirty-five percent of service members returning from the Iraq war have sought professional help for mental health issues, according to researchers at the Department of Defense. The 35 percent figure compares to 21 percent of veterans returning from Afghanistan and 24 percent of those who served elsewhere in the world. About 12 percent of the Iraq veterans were diagnosed with serious mental or emotional problems, said study author Col. Charles Hoge, M.D., of the Walter Reed Army Institute of Research.

In a separate study, released at a December 2005 Congressional hearing, Army researchers reported that 21 percent of soldiers returning from combat areas were misusing alcohol a year after their return home, although just 13 percent misused alcohol prior to deployment. The number of soldiers with anger and aggression problems increased from 11 percent pre-deployment to 22 percent post-deployment, while the divorce rate rose from 9 percent to 15 percent.

Military experts describe the Iraq war as especially stressful because it is a counterinsurgency operation with no defined front lines, meaning that soldiers rarely can relax and let down their guard.

"The most important finding is that a large number of soldiers and Marines are using mental health services very soon after they get home," Dr. Hoge noted. But some veterans' groups contend that Iraq war vets need more help than they are getting. "We've got 35 years of history from the Vietnam War that if they don't receive proper care, they turn to drugs and alcohol, they lose their homes. It's happening again," said Steve Robinson, director of the National Gulf War Resource Center.

Sources: Stars and Stripes, December 9, 2005; Hoge CW, Auchterlonie JL & Milliken CS (2006). Mental health problems, use of mental health services, and attrition from military service after returning from deployment to Iraq or Afghanistan. Journal of the American Medical Association 295:1023-1032.

Meth Bill Passes Despite White House Opposition

A bill calling for more spending on preventive advertising aimed at methamphetamine passed the House of Representatives in March on a 399-5 vote, despite a veto threat from the White House. The measure, sponsored by Rep. Mark Souder (R-IN), would require the Office of National Drug Control Policy (ONDCP) to spend at least 10 percent (\$12 million) of the funding for its national anti-drug media campaign on ads targeting methamphetamine, and to earmark another \$15 million to ads directed to "hot spots" around the country.

"The United States Congress wants some action out of this administration on meth," said Rep. Souder, who blasted the Bush administration's approach to methamphetamine as "appalling." His bill also calls for elevating the ONDCP director's post to that of a Cabinet secretary.

ONDCP spokesman Thomas Riley said the administration opposes the Sounder bill because methamphetamine is "not a problem in big cities" and that young people's use of the drug is declining. "The campaign as it is right now is working, finally, and is stretched thin," he said. "Further constraining it and further limiting it at a time when it's already very vulnerable — I worry about the effect that might have on its success."

The White House also opposed making the ONDCP directorship a Cabinet-level office, on the grounds that the bill infringes on Presidential prerogatives. Source: Join Together Online, March 12, 2006.

Drug Smuggling by Mail on the Rise, Says INCB

Use of the mails to smuggle drugs into the U.S. and other countries poses a major threat to law enforcement, says the annual report of the International Narcotics Control Board (INCB). The Board is the independent and quasi-judicial control organ that monitors implementation of the United Nations drug control conventions.

In a message accompanying the report, released March 1st in Vienna, Board President Hamid Ghodse said that, over the past five years, almost every region of the world has experienced an increase in smuggling through the mail. For instance, in the United States, where 200 billion pieces of mail are handled every year, Professor Ghodse pointed to indications that traffickers are increasingly attempting to smuggle not just illicit drugs, but also chemical precursors and pharmaceutical products through the mail. In Thailand, government agents seized half a million diazepam tablets and capsules in 12 different cases in 2004. Other seizures involved alprazolam being shipped from Thailand to the United States and clonazepam shipped from Thailand to the United Kingdom. The large quantities involved indicate that traffickers intended to distribute the drugs to the illicit market, Professor Ghodse said,

with Internet pharmacies often acting as the dealers for such shipments.

While some portion of the seized tablets are illegally obtained from licit sources (as through theft, falsified trade authorizations and individual prescriptions, pharmacies not adhering to prescription requirements, etc.), the Board's report warned that significant quantities are provided by counterfeiters. It calls counterfeiting of narcotic drugs and psychotropic substances "an important element in supplying illicit markets through illegally operating Internet pharmacies."

To stem the tide, the Board urged governments to screen incoming and outgoing international mail for drug shipments. It also recommended limiting the number of entry points for parcels to allow for a more effective control of consignments and extending screening to include the premises of international mail courier companies. And, as in all counter-trafficking activities, it called close national and international cooperation "essential." Source: International Narcotics Control Board, Annual Report, March 1, 2006; accessed at www.incb.org.

FROM THE PRESIDENT'S DESK



Dr. Elizabeth F. Howell

ASAM HONORS SCIENTIFIC, **EDUCATIONAL ACHIEVEMENTS**

Elizabeth F. Howell, M.D., FASAM

nce again this year, ASAM will honor a distinguished group of individuals who have made outstanding contributions to the field of Addiction Medicine and to the Society itself. The awardees will be honored at a gala Awards Luncheon on Saturday,

May 6th, during ASAM's Med-Sci Conference. Our 2006 awards will be presented to the following outstanding leaders:

The 2006 R. Brinkley Smithers Distinguished Scientist Award goes to Rudolf H. Moos, Ph.D., Professor of Psychiatry and Behavioral Sciences at Stanford University and Senior Research Career Scientist with the Department of Veterans Affairs. The award will be presented at the Opening Plenary Session at 9:00 a.m. Friday, May 5th. At that time, Dr. Moos will deliver the award lecture, "Common Social Factors in the Development and Remission of Alcohol Use and Other Addictive Disorders."

The John P. McGovern Award on Addiction and Society goes to Carlo C. DiClemente, Ph.D., Professor and Chair of the Department of Psychology at the University of Maryland. The McGovern Award was established in 1997 to recognize and honor an individual who has made "highly meritorious contributions to public policy, treatment, research, or prevention which has increased our understanding of the relationship of addiction and society." The award is sponsored by an endowment from the John P. McGovern Foundation.

The ASAM Annual Award for "outstanding contributions to the growth and vitality of our Society, for thoughtful leadership in the field, and for deep understanding of the art and science of addiction medicine" will be presented to James W. Smith, M.D., FASAM, and to Barry Stimmel, M.D., FASAM.


An ASAM Annual Award for "expanding the frontiers of the field of Addiction Medicine and broadening our understanding of the addiction process through research and innovation" will be presented to trauma surgeon Larry Gentilello, M.D., FACS, who has led the fight nationally to overcome statutory barriers to addiction care.

The Young Investigator Award for the best abstract submitted by an author who is within five years of receiving a doctoral degree goes to Janet Soeffing, M.D., and Hannah K. Knudsen, Ph.D.

The Medical-Scientific Program Committee Award for the abstract receiving the highest rating for scientific merit goes to Cindy Parks Thomas, Ph.D.

The Awards Luncheon is an extra-fee event. Visit the ASAM Registration Desk for tickets so that you can join us in recognizing the achievements of this distinguished group.

Detailed, practical, and clearly written...truly comprehensive in its coverage... an essential resource... Journal of the American **Medical Association** This is the guide to the science and practice of Addiction Medicine! Michael M. Miller, M.D. FASAM Medical Director, Meriter Hospital



Principles Of Addiction Third Medicine

- Completely revised and updated!
- Research-based, clinically relevant
- 1,644 pages...106 chapters
- Illustrated, indexed
- Hard cover
- Published 2003

This essential reference is available at a special price of \$175 to ASAM members and \$199 to non-members.



www.asam.org

Phone 1.800.844.8948 to Order Your Copy Today!

Visit the ASAM Publications Desk for special prices on all ASAM books...

Legislators Asked to Repeal Alcohol Exclusions

Bills to repeal UPPL and other alcohol exclusions in insurance contracts have been introduced in five states in 2006: Colorado, Connecticut, Hawaii, Illinois, and Wyoming. The status of these measures as of March 2006 follows:

Colorado. Sponsored by Rep. Angela Paccione, repeal bill H.B.06-1106 was passed by the House, introduced in the Senate and, on March 7th, was recommended for approval by the Senate Business, Labor, and Technology Committee. The Senate passed the bill with amendments March 14th: the House has voted once to concur with the Senate.

Connecticut. Introduced in late-February, S.B.425 would prohibit insurance exclusions in a state that currently has no UPPL law but does have extensive judicial precedent allowing exclusions based on intoxication. The Insurance and Real Estate Committee held a public hearing March 7th and then approved the bill March 16th. The bill has been filed with the Legislative Commissioner and could be considered by the legislature as early as the week of March 19th.

Hawaii. H.B.2401/S.B.2364, which would prohibit insurance exclusions, were introduced as part of a coordinated effort to promote a comprehensive approach to alcohol problems. An op-ed in support of the measures by Lt. Governor James Aiona was published in the Honolulu Star Bulletin March 7th. The Senate version passed and was referred to the House, where the Health Committee approved the measure March 16th and referred it to the Consumer Protection and Commerce Committee.

Illinois. Sponsored by Sen. John Cullerton, S.B.2453 was referred to the Rules Committee. where it awaits consideration.

Wyoming. H.B.7 passed through the committee process, but was defeated in a vote of the House. No further action is expected during this legislative session. Source: Ensuring Solutions to Alcohol Problems. March 17, 2006.

STUDY: UPPL Laws Still a Barrier to Care

Insurance policies too often are barriers to the use of screening and brief intervention in emergency settings, because most state governments allow insurers to sell health and accident insurance policies that will not pay for injuries that occur while the insured person is under the influence of alcohol and certain other drugs. Specifically, 35 states and the District of Columbia have so-called UPPL laws (known generically as alcohol exclusion laws, or AELs) on the books.

UPPL and other AELs create a chilling effect in hospitals that interferes with effective treatment. When benefits are denied, injured patients often cannot pay for medical care. To avoid bankrupting their patients, many physicians and hospital managers decide to avoid any activity that might result in an alcoholrelated diagnosis. As a result, one in every four U.S. trauma surgeons has had a case in which a health insurer denied coverage because a trauma victim had used alcohol or other drugs, according to a recent survey.

Study author Larry M. Gentilello, M.D., of the University of Texas Southwestern Medical School, writes that "This survey shows that the Uniform Accident and Sickness Policy Provision (UPPL) law is widely used to deny coverage across the nation...Although this law is terribly out of date, it continues to discourage surgeons from doing what they believe is best for their patients."

Eighty-two percent of the emergency physicians interviewed for Dr. Gentilello's study said they would establish alcohol screening programs if no insurance barriers existed, but only eight states have repealed their UPPL laws.

An accompanying survey of state legislators found that 89 percent believed that alcoholism is treatable, and 80 percent said offering counseling in trauma centers was a good idea. Opposition to denying insurance coverage to patients injured while under the influence of alcohol or other drugs ran at about 2 to 1 among lawmakers. "Excessive alcohol use is the leading cause of injury that we see in trauma centers and ERs across the nation, and there is broad support among both physicians and legislators for eliminating a significant barrier to diagnosis and treatment," Dr. Gentilello concluded. Sources: Gentilello LM (2005). Confronting the obstacles to screening and interventions for alcohol problems in trauma centers. Journal of Trauma, Injury Infection & Critical Care September; 59(3): S137-S143, and Gentilello LM, Samuels PN, Henningfield JE et al. (2005). Alcohol screening and intervention in trauma centers: Confidentiality concerns and legal considerations. Journal of Trauma, Injury Infection & Critical Care November; 59(5): 1250-1255.

ALCOHOL POLICY TOOLKIT AVAILABLE

To assist advocates for repeal of UPPL and other alcohol exclusion laws (AELs), Ensuring Solutions to Alcohol Problems — a policy study center at The George Washington University — has developed a toolkit filled with useful resources.

According to Eric Goplerud, Ph.D., Director of Ensuring Solutions, the toolkit (which can be downloaded from the Center's website at no charge) contains the following information:

- A state policy database contains state-specific information about alcohol exclusion laws, policies, and legislation.
- Frequently asked questions help advocates explain why an obscure insurance laws gets in the way of solving alcohol problems.
- **AEL Update** reports on the latest developments in the effort to eliminate AELs.
- A PowerPoint presentation and talking points contain quotable information about alcohol exclusions and the benefits of AEL repeal.
- Model legislation provides legislative language to prohibit alcohol exclusions.
- Sample testimony is a powerful statement in support of AEL repeal.
- An opinion piece is an essay suitable for publication in journals, newsletters, and newspapers.

To access the Alcohol Exclusions Laws Toolkit, visit Ensuring Solutions' website at WWW.ENSURINGSOLUTIONS.ORG/ RESOURCES.

Topiramate for Smoking Cessation

Alcohol misuse and smoking cause poor health, frequently co-occur, and can be difficult to treat simultaneously. A drug that can effectively treat both conditions would be of great clinical interest.

Accordingly, Bankole Johnson, M.D., Ph.D., and colleagues, who were studying whether oral topiramate improved outcomes in patients being treated for alcohol dependence, also examined whether the drug could promote smoking cessation. Using data from their previous randomized controlled trial, they compared smokers (that is, persons who smoked at least 1 cigarette a day) who had received topiramate for their alcohol use disorder (in an escalating dose of 25-300 mg per day for 12 weeks; n=45) with those who had received placebo (n=49).

Their findings suggested a clear advantage for the patients who received topiramate. At 12 weeks, 17 percent of the topiramate group and 7 percent of the placebo group reported abstinence from smoking. When the data were adjusted for potential confounders, individuals in the topiramate group were significantly more likely than those in the placebo group to report abstinence from smoking (odds ratio 4.5) and to have lower serum cotinine levels.

Although no serious adverse effects were observed, subjects in the topiramate group were significantly more likely than those in the placebo group to report paresthesias and weight loss.

Source: Johnson BA, Ait-Daoud N, Akhtar FZ et al. (2005). Use of oral topiramate to promote smoking abstinence among alcohol-dependent smokers: A randomized controlled trial. Archives of Internal Medicine 165(14):1600-1605.

St. Helena Hospital Center for Behavioral Health

Adventist Health

Napa Valley, California

Located in beautiful Napa Valley, California, St. Helena Hospital's Alcohol and Chemical Recovery Program has positions for BC, addiction medicine-certified physicians, one with Medical Director responsibilities.

This highly successful program offers 39 beds for patients who meet ASAM/DSM IV criteria. Responsibilities include pre-admission triage/assessment, direct patient care, case review and daily multi-modality team meetings. Call is after-hours weekdays and some weekends.

Competitive packages include salary, malpractice coverage, some benefits and relocation assistance.

CONTACT: Physician Recruitment, St. Helena Hospital

10 Woodland Rd., St. Helena, CA 94574

FAX: 707/963-6519

EMAIL: shhphysicianrecruiting@ah.org WEBSITE: www.sthelenahospital.org

SCREENING, BRIEF INTERVENTION IN THE SPOTLIGHT

Expanding health care providers' use of screening, brief intervention, referral and treatment (SBIRT) for alcohol and other drug problems has been designated a core strategy in the Bush administration's 2006 National Drug Control Strategy. "A key priority of this Administration has been to make drug screening and intervention programs part of the nation's existing network of health, education, law enforcement, and counseling providers," the National Drug Control Strategy declares, adding that "This effort will continue in partnership with the medical community."

Expansion of SBIRT also is the focus of a grant program recently announced by the Substance Abuse and Mental Health Services Administration (SAMHSA). To be administered by SAMHSA's Center for Substance Abuse Treatment, the SBIRT program allows only the immediate office of the Governor in each state or the highest tribal official to apply, although those offices may file a joint application with the state Substance Abuse Authority or other executive level organization.

Additional information on SBIRT and the grants program is available on SAMHSA's website (WWW.SAMHSA.GOV) . The 2006 National Drug Control Strategy can be accessed at WWW.WHITEHOUSE DRUGPOLICY.GOV.

AA Attendance **Improves Outcomes**

Alcoholics Anonymous (AA) is a preferred form of aftercare for patients discharged from formal treatment programs, but little is known about AA involvement and its effects on abstinence over time. In a study reported in the journal Alcohol: Clinical & Experimental Research, researchers assessed participation in AA, abstinence, and other alcohol outcomes over five years among 349 patients who entered treatment at baseline and attended AA at least once during the follow-up period.

The researchers described four patterns of AA attendance: "low" (mainly during the year following treatment entry); "medium" (about 60 meetings per year, with a slight increase by year 5); "high" (more than 200 meetings a year, with a slight decrease by year 5); and "declining" (almost 200 meetings the year following treatment entry and about 6 meetings in

They found that abstinence rates (past 30 days) differed significantly across the four groups. Specifically, 79 percent of patients with "high" attendance reported abstinence at year 5, followed by 73 percent with "medium" attendance, 61 percent with "declining" attendance, and 43 percent with "low" attendance. Source: Kaskutas LA, Ammon L, Delucchi K et al. (2005). Alcoholics Anonymous careers: Patterns of AA involvement five years after treatment entry. Alcohol: Clinical & Experimental Research 29(11);1983-1990.



37th Annual Medical-Scientific Conference

San Diego Sheraton – Hotel and Marina May 4-7, 2006 – San Diego, CA

- Courses and Workshops
- Abstracts and Poster Presentations
- Up to 22 Continuing Medical Education Credits
- Scientific Exhibits
- Ruth Fox Course
- Distinguished Scientist Lecture
- Buprenorphine Training Course
- Public Policy Plenary Session

















ASAM MED-SCI 2006

unny San Diego is the site of ASAM's 37th Annual Medical-Scientific Conference, where addiction experts from around the world will gather for a program rich in scientific symposia, clinical courses and workshops, and research papers and poster sessions. The conference — which welcomes ASAM members as well as non-member researchers, educators, and clinicians — is preceded on May 4th by the Ruth Fox Course for Physicians, which marks its 50th anniversary this year. The conference concludes on Sunday, May 7th, with a Buprenorphine Training Course designed to qualify ASAM members and other physicians to prescribe buprenorphine in office-based practice.

Program chair Jeffrey Samet,

M.D., M.A., M.P.H., has collaborated with co-chairs Lawrence S. Brown, Jr., M.D., M.P.H., FASAM, and Marc Galanter, M.D., FASAM, and the Program Committee to design a diverse program that affords participants an opportunity to interact with experts in the field. Major events include special day-long symposia organized by the National Institute on Alcohol Abuse and Alcoholism and the National Institute on Drug Abuse, as well as a special session on methamphetamine organized by the Center for Substance Abuse Treatment.

The Annual Business Meeting and Breakfast will be gaveled to order at 7:00 a.m. Friday, May 5th, by ASAM President Elizabeth F. Howell, M.D., FASAM. Early risers will be rewarded with a delicious buffet breakfast, to be served from 6:45 a.m., courtesy of The Christopher D. Smithers Foundation. Dr. Howell promises a highly interactive meeting, with an emphasis on soliciting members' views as to their needs and priorities, and how ASAM can best meet those needs. The breakfast and business meeting, which are open only to ASAM members, are included in the conference registration fee.



37th Medical-Scientific Conference Convenes May 4th in San Diego

The official opening of the conference, which immediately follows the business meeting, features an address by Rudolf H. Moos, Ph.D., Professor of Psychiatry and Behavioral Sciences at Stanford University and Senior Research Career Scientist with the Department of Veterans Affairs. Dr. Moos is the recipient of the 2006 R. Brinkley **Smithers Distinguished Scientist** Award. He will deliver the award lecture on "Common Social Factors in the Development and Remission of Alcohol Use and Other Addictive Disorders." Other distinguished speakers scheduled for the opening plenary are H. Westley Clark, M.D., J.D., M.P.H., FASAM, Director of the Center for Substance Abuse Treatment; Ting Kai-Li, M.D., Direc-

tor of the National Institute on Alcohol Abuse and Alcoholism; and Nora Volkow, M.D., Director of the National Institute on Drug Abuse.

The ASAM Awards Luncheon, to begin at 12:15 p.m. Saturday, May 6th, honors outstanding contributions to the addiction field, as well as those who have made notable contributions to the Society. A traditional highlight of the luncheon is the John P. McGovern Award and Lecture on Addiction and Society, established in 1997 to honor an individual who has made highly meritorious contributions to public policy, treatment, research, or prevention and who has increased our understanding of the relationship of addiction and society. The award is sponsored by an endowment from the John P. McGovern Foundation. This year's recipient is Carlo C. DiClemente, Ph.D., Professor and Chair of the Department of Psychology at the University of Maryland. (The Awards Luncheon is an extra fee event.)

Details of conference activities are found on the following pages of **ASAM News** and on ASAM's web site at WWW.ASAM.ORG.

VISITORS TO THE ASAM CONFERENCE WILL FIND MUCH TO ENJOY IN SAN DIEGO

he conference will be held at the Sheraton San Diego Hotel and Marina, 1380 Harbor Island Drive, San Diego, CA 92101. Hotel features include outdoor pools, fitness and spa facilities, high-speed Internet access in all guest rooms, concierge service, and beaches nearby. The Sheraton also provides complimentary airport shuttle service, with vans operating every 20-25 minutes. Self-parking is available at the hotel for \$16 a night; valet parking is \$22 per day.

A limited number of guest rooms are available at the special conference rate of \$187 single or double. Rates are subject to state, local and occupancy taxes, currently 15% (subject to change). To make your reservation, contact the Sheraton San Diego Reservations Department directly at 619/291-2900 or use the Sheraton's central reservation line at 1-800/325-3535. All reservations must be accompanied by a deposit equal to one night's room and tax, or be

guaranteed with a major credit card number and expiration date. In order to receive the conference rate, tell the reservation agent that you are attending the "ASAM Medical Scientific Conference." The hotel cannot guarantee availability of rooms at the conference rate after April 10th, so make your reservations early!

Although the Med-Sci Conference offers more than enough activities to keep registrants busy, the conference hotel site offers great opportunities to explore San Diego, where the average year-round temperature is 72 degrees. The Sheraton is situated on Harbor Island in beautiful San Diego Bay, and offers panoramic views of the bay and the downtown city skyline. Unique attractions within a 10-minute drive of the hotel include Sea World, the San Diego Zoo, historic Old Town San Diego, Balboa Park, and Seaport Village. For more information, visit the hotel's website at WWW.SHERATON.COM/ SANDIEGOMARINA.

CONFERENCE REGISTRATION AND FEES

Register on-site at the ASAM Conference Registration and Information Desk, which will be open during the following hours:

Wednesday, May 3rd 5:00 p.m. to 8:00 p.m. Thursday, May 4th 6:30 a.m. to 6:30 p.m. Friday, May 5th 7:00 a.m. to 6:00 p.m. Saturday, May 6th 7:00 a.m. to 6:00 p.m. Sunday, May 7th 7:00 a.m. to 1:00 p.m.

The following fees apply to on-site registrations only:

Ruth Fox Course for Physicians

(Thursday, May 4th) ASAM member\$200 Non-member physician\$250 Non-physician professional (R.N., Ph.D., CAC, LCSW, etc.) \$175 Resident, Fellow, Intern (with proof of status)\$150 Student (with proof of status)\$75

37th Annual Medical-Scientific Conference

(May 5-7th) ASAM member \$475 Non-member physician\$585 Non-physician professional (R.N., Ph.D., CAC, LCSW, etc.) \$475 Paper presenter\$350 Resident, Fellow, Intern (with proof of status) No charge Student (with proof of status) No charge

Daily Registration for the Medical-Scientific Conference

.....\$225 per day

ASAM Awards Luncheon

(Saturday, May 6th, 1:00 p.m.)\$50 for Medical-Scientific Conference registrants

Buprenorphine and Office-Based Treatment of Opiod Dependence

(Sunday, May 7th) ASAM member\$100 Non-member physician\$125 Non-physician professional (R.N., Ph.D., CAC, LCSW, etc.) \$75

CONFERENCE PROGRAM COMMITTEE

Chair: Jeffrey Samet, M.D., M.A., M.P.H. Co-Chair: Lawrence S. Brown Jr., M.D., M.P.H., FASAM

Co-Chair: Marc Galanter, M.D., FASAM

Members:

Gregory C. Bunt, M.D. David A. Fiellin, M.D. Mark S. Gold, M.D. R. Jeffrey Goldsmith, M.D. Enoch Gordis, M.D. David A. Gorelick, M.D., Ph.D. Edward Gottheil, M.D., Ph.D. James A. Halikas, M.D., FASAM Jag H. Khalsa, Ph.D. Donald J. Kurth, M.D., FASAM Norman S. Miller, M.D., FASAM Alfonso Paredes, M.D. Richard K. Ries, M.D., FASAM Peter D. Rogers, M.D., M.P.H., FASAM Barry Stimmel, M.D., FASAM Anton Che' Bizzell, M.D., CSAT Liaison Dorynne Czechowicz, M.D., NIDA Liaison Carlton K. Erickson, Ph.D., RSA Liaison Norman W. Wetterau, M.D., AAFP Liaison Mark Willenbring, M.D., NIAAA Liaison Joan Ellen Zweben, Ph.D., Consultant

Staff:

Eileen McGrath, J.D., Executive Vice President/CEO Sandy S. Metcalfe, Director of Meetings and Conferences Berit Boegli, Exhibit Program Coordinator

MEET YOUR REGIONAL DIRECTOR

During the Dessert Reception on Friday, May 5th, from 9:00 to 11:00 pm, members will have an opportunity to meet informally with **ASAM's Regional Directors** and discuss issues of importance to their Chapters and Regions.

Ruth Fox Course for Physicians Marks Its 25th Anniversary

Join your colleagues as ASAM celebrates the 25th Anniversary of the Ruth Fox Course for Physicians on May 4th from 8:00 a.m. to 5:30 pm. The course is dedicated to providing practicing physicians with cutting-edge knowledge about current trends in the field of addiction medicine.

The first Ruth Fox course was organized by the late Maxwell N. Weisman, M.D., and held May 5, 1980, in Seattle, Washington, in conjunction with the National Alcoholism Forum. Over the ensuing 25 years, the Ruth Fox Course has educated thousands of physicians.

This year's course, which is co-chaired by Doctors Margaret A.E. Jarvis, Louis E. Baxter, Sr., and John C. Tanner, features an address by John P. Walters, Director of the Office of National Drug Control Policy, as well as the following outstanding sessions:

- Remembering Dr. Ruth Fox Looking Back/Looking Forward: Stanley E. Gitlow, M.D., FACP, FASAM
- Origins of ASAM/Addiction Medicine: David C. Lewis, M.D.
- Spirituality: A Tool for the Clinician's Armamentarium: John N. Chappel, M.D., FASAM
- Alcohol Screening & Interventions in Trauma & Emergency Departments: Larry M. Gentilello, M.D., FACS
- Professionalism: Boundaries, Principles & Ethics in Addiction Practice: James C. Montgomery, M.D.
- Impaired and Disruptive Physicians: Peter A. Mansky, M.D.
- What Influences the "Tipping Point" in the U.S. or International Tobacco **Epidemic?** Lynda Hyder Ferry, M.D.
- Recovery in the Middle East: Keith Humphreys, Ph.D.
- Treatment in the Criminal Justice System: Jeffrey D. Baxter, M.D.
- Literature Review: David R. Gastfriend, M.D.

QUICK GUIDE TO MED-SCI PROGRAM EVENTS

THURSDAY, MAY 4, 2006

6:00 - 8:00 pm

Welcome Reception and Opening of the **ASAM Exhibit Hall**

8:00 - 10:00 pm

COMPONENT SESSION I: Opioid Agonist Treatment 2005 — Training, Clinical **Practice and Policy Initiatives**

Sponsored by the ASAM Sub-Work Groups on Opioid Agonist Treatment and **Buprenorphine Training**

SPEAKERS: Daniel P. Alford, M.D., M.P.H., David A. Fiellin, M.D., Judith Martin, M.D., Edwin Salsitz, M.D., FASAM, Deb Stephenson, M.D.. & Richard A. Yoast, Ph.D.

COMPONENT SESSION II: Why Not Private Practice? Practice Management: Coding and Billing Strategies

CHAIRS: A. Kenison Roy III, M.D., FASAM & John P. Femino, M.D., FASAM

COMPONENT SESSION III: Public Policy in Addiction Medicine: Past, Present, and Future

Sponsored by the ASAM Public Policy Committee

SPEAKERS: Mark L. Kraus, M.D., FASAM & Petros Levounis, M.D., M.A.

COMPONENT SESSION IV: Addiction Medicine and Primary Care

SPEAKERS: Bonnie B. Wilford, M.S. & Norman Wetterau, M.D., FAAFP

COMPONENT SESSION V: Chapters Council Sponsored by the Chapters Council and the ASAM State Medical Specialty Society Program

FRIDAY, MAY 5, 2006

7:00 am

ASAM Annual Business Meeting & Breakfast

(ASAM Members only —breakfast service will start at 6:45 am)

Continental Breakfast for ASAM Guests and Non-Members

8:30 am

Opening Scientific Plenary and R. Brinkley **Smithers Distinguished Scientist Lecture**

"Common Social Factors in the Development and Remission of Alcohol Use and Other Addictive Disorders"

SPEAKER: Rudolph H. Moos, Ph.D., Senior Research Career Scientist, Department of Veterans Affairs, and Professor of Psychiatry and Behavioral Sciences, Stanford University, Menlo Park, CA

ASAM Exhibit Hall Opens

10:00 am

Refreshment Break — ASAM Exhibit Hall

10:30 am - 12:30 pm and 2:30 pm - 6:00 pm

SYMPOSIUM 1: Clinical Approaches to HIV and Hepatitis C Infections in **Drug Abuse**

Sponsored by the National Institute on Drug Abuse (NIDA)

ORGANIZERS: Jag H. Khalsa, Ph.D. & Frank J. Vocci, Jr., Ph.D.

SPEAKERS: Andrea Cox, M.D., Ph.D., Brian Edlin, M.D., M.P.H., Richard Garfein, Ph.D., M.P.H., Ramesh Ganju, Ph.D., Kristy Marie Hendricks, Sc.D., Charles B. Hinkin, Ph.D., ABPP, Robert Horsburgh, M.D., Kenneth E. Sherman, M.D., Ph.D., Jack Stapleton, M.D., Ellen Tedaldi, M.D. & Terry Wright, M.D.

10:30 am - 12:30 pm

SYMPOSIUM 2: Crossing the Quality **Chasm in Addiction Medicine**

SPEAKERS: A. Thomas McLellan, M.D., Jeffrey Samet, M.D., M.A., M.P.H. & Constance Weisner, Dr.P.H., M.S.W.

COURSE 1: Criminal Justice System and the Therapeutic Community: Treatment Alternatives to Incarceration

SPEAKERS: George DeLeon, Ph.D., Terry Horton, M.D., Donald J. Kurth, M.D., FASAM, Richard Rosner, M.D. & Anne J. Swern

WORKSHOP A: Drug Seeking or Pain? **Teaching About Addiction and Chronic Pain**

SPEAKERS: Karen Cropsey, Psy.D., Laura Morgan, Pharm.D. & Michael Weaver, M.D., FASAM

WORKSHOP B: Treatment of Methamphetamine Dependence in a Residential **Drug Court Therapeutic Community:** A Novel Pilot Project

SPEAKERS: John B. Averitt, Ph.D., Roland Gray, M.D., Seth Norman, J.D. & David K. Patzer, M.D.

10:30 am - 1:00 pm **PAPER SESSION 1**

SAVE MONEY!

Attend the Medical-Scientific Conference at the member rate by joining ASAM now!

12:30 pm - 2:00 pm

Lunch break

1:00 - 2:00 pm

POSTER SESSION

2:00 pm - 4:00 pm

SYMPOSIUM 4: Integrated Treatment of the Dually Diagnosed

ORGANIZER: Marc Galanter, M.D., FASAM SPEAKERS: Petros Levounis, M.D., Jaime Grodzicki, M.D. & Richard K. Ries, M.D., **FASAM**

2:00 pm - 4:30 pm **PAPER SESSION 2**

2:00 pm - 6:00 pm

SYMPOSIUM 1 (continued): Clinical Approaches to HIV and Hepatitis C Infections in Drug Abusers

SYMPOSIUM 3: Treating the Physician — **Engagement, Treatment Approaches** and Outcomes

ORGANIZER: Peter A. Mansky, M.D. SPEAKERS: Louis E. Baxter, Sr., M.D., FASAM, Lynn R. Hankes, M.D., FASAM & Margaret A E. Jarvis, M.D., FASAM

2:30 pm - 4:30 pm

COURSE 2: The Psychotherapy and Treatment of Methamphetamine Addiction SPEAKERS: Steven J. Lee. M.D. & Petros Levounis, M.D., M.A.

WORKSHOP C: Addiction in Pregnancy: **Motivational Enhancement Treatment**

SPEAKER: James J. Nocon, M.D.

6:30 - 8:30 pm

Ruth Fox Endowment Reception (by invitation only)

8:00 - 10:00 pm

WORKSHOP D: Hospital Based Substance Abuse Consultation Services — Clinical and Administrative Experiences

SPEAKERS: Rita Azalos, M.D., Richard Blondell, M.D., Michael Weaver, M.D. & Christopher Welsh, M.D.

WORKSHOP E: Group Therapy of **Substance Abuse**

SPEAKERS: David W. Brook, M.D., FASAM & Marc Galanter, M.D., FASAM

WORKSHOP F: Narcotics Anonymous (NA) — A Vital Resource in the Treatment of Addiction

SPEAKERS: Bob MacFarlane, M.D., Donna Markus, Ph.D. & Bob Stewart

9:30 pm

ASAM Dessert and Coffee Reception (All registrants are invited)

SATURDAY, MAY 6, 2006

7:00 am - 8:00 am Continental Breakfast — **ASAM Exhibit Hall**

8:00 am - 9:30 am

PUBLIC POLICY PLENARY: The 30 Patient Rule — Rational, Irrational, or Both?

Sponsored by ASAM's Public Policy and Legislative Advocacy Committees Invited Panel Members: CSAT Director H. Westley Clark, M.D., JD, M.P.H., FASAM, Mark L. Kraus, M.D., FASAM, Petros Levounis, M.D., M.A., and DEA Administrator Karen Tandy

9.30 am

Refreshment Break - ASAM Exhibit Hall

10:00 am - 12:00 Noon and 2:30 pm - 5:30 pm

SYMPOSIUM 5: Medications Development for Alcoholism: From the Bench to the Patient

Sponsored by the National Institute on Alcohol Abuse and Alcoholism (NIAAA) ORGANIZERS: Mark L. Willenbring, M.D. & Raye Z. Litten, Ph.D.

SPEAKERS: Raymond F. Anton, M.D., Jack R. Cornelius, M.D., M.P.H., Bankole A. Johnson, M.D., Ph.D., George F. Koob, Ph.D., Henry R. Kranzler, M.D., John M. Littleton, M.D., Ph.D., Barbara J. Mason, Ph.D. & Mark L. Willenbring, M.D.


SYMPOSIUM 6: Health Care Disparities, **Quality of Care, and Pay for Performance:** What Is An Addiction Medicine Specialist to Know and Do?

ORGANIZER: Lawrence S. Brown, Jr., M.D., M.P.H., FASAM

SPEAKERS: John C. Nelson, M.D., M.P.H., Randall W. Maxey, M.D., Ph.D., and Matthew Wynia, M.D.

COURSE 3: Teaching Addiction Medicine to Medical Students, Residents and **Primary Care Providers**

SPEAKERS: R. Jeffrey Goldsmith, M.D. & Edwin A. Salsitz, M.D.



Balboa Park in San Diego

WORKSHOP G: Al-Anon/Alateen **Presents Recovery From The Family Disease of Addiction**

ORGANIZER: Joseph A. Troncale, M.D. SPEAKERS: Neil A. Capretto, D.O., FASAM, Claire Ricewasser & Jeffrey D. Roth, M.D.,

WORKSHOP H: Crystal, Coke, Sex and the Net: Treating the Dangerous and **Difficult Combinations**

SPEAKERS: M. Deborah Corley, Ph.D. & James C. Montgomerv, M.D.

WORKSHOP I: Monitoring Alcohol **Abstinence: Using EtG and Other Markers** and Devices

ORGANIZER: Gregory E. Skipper, M.D. SPEAKERS: Michael Feldman, Ph.D.,

Michel Sucher, M.D., Douglas Stellato Kabat, L.C.S.W. & Friedrich Wurst, M.D.

12:15 pm

ASAM Awards Luncheon

2:30 - 4:30 pm

SYMPOSIUM 7: Lost in Translation: Breaking the Bottleneck of Research Adoption

ORGANIZER: Joan E. Zweben, Ph.D. SPEAKERS: Jeanne L. Obert, M.A., M.S.N., Jack B. Stein, Ph.D.& Mark L. Willenbring, M.D.

SYMPOSIUM 8: Rural Addiction Medicine: Real Need and Special Opportunity ORGANIZER: Noman Wetterau, M.D., FAAFP, FASAM

SPEAKERS: Marcello A. Maviglia, M.D., M.P.H., Berton Toews, M.D. & Art Van Zee, M.D.

COURSE 4: Treatment Approaches for Persons in the Criminal Justice System Who Have Co-occurring Substance Use and Mental Disorders

SPEAKERS: Charlene Le Fauve, Ph.D., Fred C. Osher, M.D., & Roger H. Peters, Ph.D.

WORKSHOP J: Buprenorphine on the Cutting Edge: Advanced Issues in the Clinical Use of Buprenorphine

SPEAKERS: Eric D. Collins, M.D., Hendree Jones, Ph.D., Edwin A. Salsitz, M.D., FASAM & Michael Weaver, M.D., FASAM

Refreshment Break - ASAM Exhibit Hall

6:00 - 9:00 pm

DINNER AND CME COURSE: Advances in Pharmacotherapy for Alcohol Dependence

SPEAKERS: Charles O'Brien, M.D., Ph.D., David Mee-Lee, M.D., Gerald Shulman, M.A., FACATA & Marc J. Fishman, M.D. (There is no charge for dinner, but registration is required)

SUNDAY, MAY 7, 2006

8:00 am

Continental Breakfast

8:30 - 10:30 am

SYMPOSIUM 11: Pathological Gambling: A Non-Substance, Substance Use Disorder? ORGANIZER: Edward Gottheil, M.D., Ph.D.

SPEAKERS: Nady el-Guebaly, M.D., Jon E. Grant, J.D., M.D., Clayton Neighbors, Ph.D. & Ken C. Winters, Ph.D.

COURSE 5: Six 12 Step-Oriented **Principles in Addiction Treatment**

SPEAKER: Daniel H. Angres, M.D.

WORKSHOP K: Addiction and Sleep Disorders

SPEAKERS: Jeff Daiter, M.D., CCFP, FCFP & Michael Varenbut, M.D., CCFP, FCFP

WORKSHOP L: Physician Health Programs — Effective Linking of Monitoring, 12-Step Fellowship Participation and Addiction Treatment — A Possible New Treatment Paradigm

ORGANIZER: Robert L. DuPont, M.D., **FASAM**

SPEAKERS: A. Thomas McLellan, Ph.D., Gregory E. Skipper, M.D. & William L. White, M.A.

WORKSHOP M: Addiction Treatment Using Spirituality Enhanced Twelve Step Facilitation (TSF) and Network Therapy in **Residential Treatment for Physicians and** Other Professionals

SPEAKER: John S. Sappington, M.D., FASAM

WORKSHOP N: Watching Over A Shoulder: Measurable Monitoring of Behavioral Addictions in the Workplace

SPEAKERS: M. Deborah Corley, Ph.D. & James C. Montgomery, M.D.

8:30 am - 12:30 pm

SYMPOSIUM 9: Treatment of Methamphetamine Use Disorders Sponsored by the Center for Substance Abuse Treatment (CSAT)

ORGANIZER: Anton Ché Bizzell, M.D. SPEAKERS: H. Westlev Clark, M.D., J.D., M.P.H., FASAM, Rachel Gonzales, Ph.D., William Haning, M.D., Walter Ling, M.D., Jane C. Maxwell, Ph.D., Thomas F. Newton, M.D., Martin Paulus, M.D., Richard Rawson, Ph.D. & Matt Torrington, M.D.

10:30 am

Refreshment Break

11:00 am - 1:00 pm

WORKSHOP O: Addiction Treatment in Correctional Settings

SPEAKERS: Jeffrey Baxter, M.D. & Stacy Seikel, M.D.

End of Conference

Older Americans Misusing Alcohol, Opiates

More older Americans are entering addiction treatment, primarily because of problems with alcohol and prescription opiates, according to data released by the Substance Abuse and Mental Health Services Administration (SAMHSA).

Data collected for the 2003 Treatment Episode Data Set (TEDS) show that treatment admissions among Americans ages 55 and older increased by 32 percent between 1995 and 2002. While alcohol remains the primary drug of abuse among older adults, a growing number cited prescription opiates: 12 percent in 2002, compared to 6.8 percent in 1995. Only 4 percent reported cocaine abuse, 3 percent reported marijuana abuse, and 1 percent reported abuse of stimulants. All of the rates were lower than those for the general population, except that the reports of alcohol abuse were higher among seniors than among patients of all ages.

Explaining that alcohol and other drug problems often are overlooked or misdiagnosed in older adults, SAMHSA Administrator Charles Curie said, "Too often, family members are ashamed of the problem and choose not to address it. Health care providers tend not to ask older patients about alcohol abuse.... Sometimes, the symptoms are mistaken for those of dementia, depression, or other problems common to older adults."

To counter the upward trend in abuse of opiates, SAMHSA and the Food and Drug Administration (FDA) have launched a public education campaign to encourage older adults to "Do The Right Dose" when using prescription pain relievers. The campaign will strive to educate older adults that prescription pain medications are safe and effective when used correctly, but if misused, could lead to addiction or other problems. The educational materials can be accessed online at HTTP:// ASYOUAGE.SAMHSA.GOV/DOTHERIGHTDOSE/.


Free Slideshow CD on **Using the NIAAA** Clinician's Guide

NIAAA presents a new instructional CD for using Helping Patients Who Drink Too Much - A Clinician's Guide. The CD features:

- A slideshow overview of the Guide with step-by-step instructions
- · Slide templates and animated graphics ideal for customizing presentations about the Guide

Download the Guide and related tools at www.niaaa.nih.gov





Surgeon General Issues Call to Action on Underage Drinking



Surgeon General Richard Carmona, M.D., M.P.H.

National attention to the problem of youth alcohol use picked up momentum last month as Surgeon General Richard Carmona, M.D., M.P.H., officially announced his intention to issue a Call to Action on underage drinking.

At a December 2005 meeting convened by the federal government's **Interagency Coordinating**


Committee on the Prevention of Underage Drinking (ICCPUD), Dr. Carmona promised that he would use his "bully pulpit" to address underage drinking. He followed through in February 2006, when his office issued a request for comments on the proposed Call to Action, with a deadline of March 15th for public input.

A 2003 report from the National Academy of Sciences, "Reducing Underage Drinking: A Collective Responsibility," is widely viewed as a model for the Surgeon General's initiative. The Congressionally mandated NAS report, issued by the National Research Council and the Institute of Medicine, advised the federal government to educate adults about existing laws and the consequences of underage drinking, urged parents to do a better job supervising their children, and called on the alcohol and entertainment industries to shield youth from unsuitable messages about alcohol consumption.

To fund the proposed activities, the report urged that Congress and state legislatures raise alcohol excise taxes — particularly on beer, which studies show is the alcoholic beverage that most young people prefer. Other NAS recommendations included increased compliance checks on retailers, backed by the threat of states losing federal funding if enforcement falls short.

Many observers doubt that the Bush administration will confront issues like alcohol taxes and advertising, and Dr. Carmona is barred from making specific policy recommendations as part of his Call to Action. On the other hand, Ohio First Lady Hope Taft notes that former Surgeon General Everett Koop's landmark report on tobacco didn't include policy recommendations, either, but still led to a paradigm shift in how the public viewed smoking. "We hope that this is the beginning of a shift on childhood drinking," she told Join Together.

The Surgeon General's report is expected to be released in April to correspond with Alcohol Awareness Month. Source: Bob Curley, Join Together Online, March 13, 2006.



• 2 to 3 times more patients maintained abstinence vs placebo in long- and short-term studies, respectively²

- Works well with a variety of psychosocial therapies³⁻⁶
- Excellent safety and tolerability profile¹⁻⁷
- Unique mechanism of action is thought to restore neurotransmitter balance*1
- Used in over 1.5 million patients worldwide⁷

CAMPRAL® (acamprosate calcium) is contraindicated in patients with severe renal impairment (creatinine clearance ≤30 mL/min). CAMPRAL is contraindicated in patients with known hypersensitivity to acamprosate calcium or any excipients used in the formulation. CAMPRAL does not eliminate or diminish withdrawal symptoms. Alcohol-dependent patients, including those patients being treated with CAMPRAL, should be monitored for the development of symptoms of depression or suicidal thinking. The most common adverse events reported with CAMPRAL vs placebo (≥3% and higher than placebo) were asthenia, diarrhea, flatulence, nausea, and pruritus.

*The mechanism of action of acamprosate in the maintenance of abstinence is not completely understood. Chronic alcohol exposure is hypothesized to alter the normal balance between neuronal excitation and inhibition. In vitro and in vivo studies in animals have provided evidence to suggest acamprosate may interact with neurotransmitter systems centrally, and has led to the hypothesis that acamprosate restores this balance. The clinical significance in humans is unknown.

References: 1. CAMPRAL® (acamprosate calcium) Delayed-Release Tablets Prescribing Information, Forest Laboratories, Inc., St Louis, Mo, 2004. 2. Data on file, Forest Laboratories, Inc. 3. Pelc I, Verbanck P, Le Bon O, Gavrilovic M, Lion K, Lehert P. Efficacy and safety of acamprosate in the treatment of detoxified alcohol-dependent patients: a 90-day placebo-controlled dose-finding study. Br J Psychiatry. 1997;17:73–77. 4. Sass H, Soyka M, Mann K, Zieglgansberger W. Relapse prevention by acamprosate: results from a placebo-controlled study on alcohol dependence. Arch Gen Psychiatry. 1996;53:673–680. 5. Paille FM, Guefi JD, Perkins AC, Royer RJ, Steru L, Parot P. Double-blind randomized multicentre trial of acamprosate in maintaining abstinence from alcohol. Alcohol Alcohol. 1995;30:239–247. 6. Pelc I, Ansons C, Lehert P, et al. The European NEAT Program integrated approach using acamprosate and psychosocial support for the prevention of relapse in alcohol-dependent patients with a statistical modeling of therapy success prediction. Alcohol. 2003;16:203–215.

Please see Brief Summary of Prescribing Information on the following page.

CAMPRAL is a registered trademark of Merck Santé s.a.s., subsidiary of Merck KGaA, Darmstadt, Germany





Brief Sun

Brief Summary: For complete details, please see full Prescribing Information for CAMPRAL

INDICATIONS AND USAGE

CAMPRAL (acamprosate calcium) is indicated for the maintenance of abstinence from alcohol in patients with alcohol dependence who are abstinent at treatment initiation. Treatment with CAMPRAL should be part of a comaround dependence who are assistant at dearniest initiation. Treatment with CAWIFPAL stroug depart of a comprehensive management program that includes psychosocial support. The efficacy of CAMIFPAL in promoting abstinence has not been demonstrated in subjects who have not undergone detoxification and not achieved alcohol abstinence prior to beginning CAMIPPAL treatment. The efficacy of CAMIPPAL in promoting abstinence from alcohol in polysubstance abusers has not been adequately assessed.

CONTRAINDICATIONS

CAMPRAL is contraindicated in patients who previously have exhibited hypersensitivity to acamprosate calcium or any of its components. CAMPRAL is contraindicated in patients with severe renal impairment (creatinine clearance ≤30 mL/min).

CAMPRAL does not eliminate or diminish withdrawal symptoms. General: Renal Impairment Treatment with CAMPRAL in patients with sweer ernal impairment (creatinine clearance of 30-50 ml/min) requires a dose reduction. Patients with severe renal impairment (creatinine clearance of 30-50 ml/min) requires a dose reduction. Patients with severe renal impairment (creatinine clearance of 30-50 ml/min) should not be given CAMPRAL, (see also CONTRAINDICATIONS). Suicidaility in controlled clinical trials of CAMPRAL, adverse events of a suicidal nature (suicidal ideation, suicide attempts, completed suicides) were infrequent overall, but were more common in CAMPRAL redated patients than in patients treated with placebo (1.4% vs. 0.5% in studies of 6 months or less; 2.4% vs. 0.8% in year-long studies). Completed suicides occurred in 3 of 2272 (0.13%) patients in the pooled acamprosate group from all controlled studies and 2 of 1962 patients (0.10%) in the placebo group. Adverse events coded as "depression" were reported at similar rates in CAMPRAL-treated and placebo-treated patients. Although many of these events occurred in the context of alcohol relapse, no consistent pattern of relationship between the clinical course of recovery from alcoholism and the emergence of suicidality was identified. The interrelationship between alcohol dependence, depression and suicidality is well-recorpized and complex. Alcohol-dependent patients, including those patients being treated with CAMPRAL, should be anothered for the development of symptoms of depression or suicidal thinking. Families and caregivers of patients being treated with CAMPRAL should be alerted to the need to monitor patients for the emergence of symptoms of depression or suicidality, and to report such symptoms to the patient's health care provider. Information for Patients Physicians are advised to discuss the following issues with patients for thom they prescribe CAMPRAL Any psychoactive drug may impair judgment, thinking, or motor skills. Patients should be cautioned about operating hazardous machinery, including automobiles, until the patients should be activ ose of otwer the construction of the construc when it given in toxes that are approximately 3 times the human dose (on a mg/m² basis). Acamprosate calcium produced a dose-related increase in the number of fetuses with malformations in rats at oral doses of 300 mg/kg/day or greater (approximately equal to the maximum recommended human daily oral dose on a mg/m² basis). The malformations included hydronephrosis, malformed iris, retinal dysplasia, and retroesophageal subclavian artery. No findings were observed at an oral dose of 50 mg/kg/day (approximately one-fifth the maximum recommended human daily oral dose on a mg/m² basis). An increased incidence of hydronephrosis was also noted in Burgundy Tawny rabbits at oral doses of 400 mg/kg/day or greater (approximately 3 times the maximum recommended human daily oral dose on a mg/m² basis). An dovelopmental effects were observed in Nev Zealand white rabbits at oral doses up to 1000 mg/kg/day (approximately 8 times the maximum recommended human daily oral dose on a mg/m² basis). The findings in animals should be considered in relation to known adverse developmental effects of ethyl alcohol, which include the characteristics of fetal alcohol syndrome (cranifactial dysmorphism, intrauterine and postnatal growth retardation, retarded psychomotor and intellectual development) and milder forms of neurological and behavioral disorders in humans. There are no adequate and well controlled studies in pregnant women. CAMPRAL should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. **Nonteratogenic Effects** A study conducted in pregnant mice that were administered acamprosate calcium by the oral route starting on Day 15 of gestation through the end of lactation on postnatal day 28 demonstrated an increased incidence of still-born fetuses at doses of 960 mg/kg/day or greater (approximately 2 times the maximum recommended human daily oral dose on a mg/m² basis). Labor and Delivery The potential for CAMPRAL to affect the duration of labor and delivery is union should be exercise

AND ADMINISTRATION). ADVERSE REACTIONS

ADVERSE REACTIONS

The adverse event data described below reflect the safety experience in over 7000 patients exposed to CAMPRAL for up to one year, including over 2000 CAMPRAL exposed patients who participated in placebo-controlled trials. Adverse Events Leading to Discontinuation in placebo-controlled trials of 6 months or less, 8% of CAMPRAL-treated patients discontinuation in placebo-controlled trials of 6 months or less, 8% of CAMPRAL-treated patients longer than 6 months, the discontinuation rate due to adverse events was 7% in both the placebo-treated and the CAMPRAL-treated patients. Only diarrhea was associated with the discontinuation of more than 1% of patients (2% of CAMPRAL-treated patients). Only diarrhea was associated with the discontinuation of more than 1% of patients, were nevertheless more commonly cited in association with discontinuation in less than 1% of patients, were neverthed patients. Common Adverse Events Reported in Controlled Trials Common, non-serious adverse events were collected spontaneously in some controlled studies and using a checklist in other studies. The overall profile of adverse events was similar using either method. Table 1 shows those events that occurred in any CAMPRAL

treatment group at a rate of 3% or greater and greater than the placebo group in controlled clinical trials with spontaneously reported adverse events. The reported frequencies of adverse events represent the proportion of individuals who experienced, at least once, a treatment-emergent adverse event of the type listed, without regard to the causal relationship of the events to the drug.

Table 1. Events Occurring at a Rate of at Least 3% and Greater than Placebo in any CAMPRAL Treatment Group in Controlled Clinical Trials with S

Body System/ Preferred Term	CAMPRAL 1332 mg/day	CAMPRAL 1998 mg/day ¹	CAMPRAL Pooled ²	Placebo	
Number of Patients in 397 Treatment Group		1539	2019	1706	
Number (%) of Patients with an AE	248 (62%)	910(59%)	1231 (61%)	955 (56%)	
Body as a Whole	121 (30%)	513(33%)	685 (34%)	517(30%)	
Accidental Injury*	17 (4%)	44 (3%)	70 (3%)	52 (3%)	
Asthenia	29 (7%)	79 (5%)	114(6%)	93 (5%)	
Pain	6 (2%)	56 (4%)	65 (3%)	55 (3%)	
Digestive System	85 (21%)	440(29%)	574 (28%)	344(20%)	
Anorexia	20 (5%)	35 (2%)	57 (3%)	44 (3%)	
Diarrhea	39 (10%)	257(17%)	329(16%)	166(10%)	
Flatulence	4 (1%)	55 (4%)	63 (3%)	28 (2%)	
Nausea	11 (3%)	69 (4%)	87 (4%)	58 (3%)	
Nervous System	150(38%)	417(27%)	598 (30%)	500(29%)	
Anxiety**	32 (8%)	80 (5%)	118(6%)	98 (6%)	
Depression	33 (8%)	63 (4%)	102(5%)	87 (5%)	
Dizziness	15 (4%)	49 (3%)	67 (3%)	44 (3%)	
Dry mouth	13 (3%)	23 (1%)	36 (2%)	28 (2%)	
Insomnia	34 (9%)	94 (6%)	137(7%)	121(7%)	
Paresthesia	11 (3%)	29 (2%)	40 (2%)	34 (2%)	
Skin and Appendages	26 (7%)	150(10%)	187 (9%)	169(10%)	
Pruritus	12 (3%)	68 (4%)	82 (4%)	58 (3%)	
Sweating	11 (3%)	27 (2%)	40 (2%)	30 (2%)	

*includes events coded as "fracture" by sponsor; **includes events coded as "nervoursess" by sponsor
i includes 258 patients treated with acamprosate calcium 2000 mg/day, using a different dosage strength and
regimen. 2 includes all patients in the first two columns as well as 83 patients treated with acamprosate calcium 3000 mg/day, using a different dosage strength and regimen.

Other Events Observed During the Premarketing Evaluation of CAMPRAL
Following is a list of terms that reflect treatment-emergent adverse events reported by patients treated with
CAMPRAL in 20 clinical trials (4461 patients treated with CAMPRAL, 3526 of whom received the maximum
recommended dose of 1998 mg/day for up to one year in duration). This listing does not include those events
already listed above; events for which a drug cause was considered remote; event terms which were so general as
to be uninformative; and events reported only once which were not likely to be acutely life-threatening. already listed above; events for which a drug cause was considered remote; event terms which were so general as to be uninformative; and events reported only once which were not likely to be acutely life-threatening. Events are further categorized by body system and listed in order of decreasing frequency according to the following definitions: frequent adverse events are those occurring in at least 17.100 patients (only those not already listed in the summary of adverse events in controlled trials appear in this listing); infrequent adverse events are those occurring in 17.100 to 17.1000 patients. Body as a Whole — Frequent: Pendadche, abdominal pain, back pain, infection, flu syndrome, chest pain, chills, suicide attempt; Infrequent: fever, intentional overdose, malaise, allergic reaction, abscess, neck pain, hernia, intentional injury, Pare: ascites, face edema, photosensitivity reaction, abdomen enlarged, sudden death. Cardiovascular System — Frequent: palpitation, syncope; Infrequent: hypotension, tachycardia, hemorrhage, angina pectoris, migraine, varicose vein, myocardial infarct, phlebitis, postural hypotension; Rare: heart failure, mesenteric arterial occlusion, cardiomyopathy, deep thrombophlebitis, shock. Digestive System — Frequent: worniting, dyspepsia, constipation, increased appetite; Infrequent: liver function tests abnormal, gastroenteritis, gastritis, dysphagia, eructation, gastrointestinal hemorrhage, pancreatitis, rectal hemorrhage, liver cirrhosis, esophagitis, dysphagia, eructation, gastrointestinal hemorrhage, pancreatitis, rectal hemorrhage, liver cirrhosis, duodenal ulcer, mouth ulceration, carcinoma of liver. Endocrine System — Rare: goiter, hypothyroidism. Hemic and Lymphatic System — Infrequent: anemia, ecchymosis, eosinophilia, lymphocytosis, thrombocytopenia; Rare: leukopenia, lymphadenopathy, monocytosis. Metabolic and Nutritional Disorders — Frequent: hypothyroidism. Hemic increased, hyponatremia, lactic dehydrogenase increased. Musculoskeletal System — Frequent: myalgia, arthral epistaxis, pneumonia; Rare: laryngismus, pulmonary embolus. **Skin and Appendages**—Frequent: rash; infrequent: acne, eczema, alopecia, maculopapular rash, dry skin, urticaria, exfoliative dermatitis, vesiculobullous rash; Rare: psoriasis. **Special Senses**—Frequent: abnormal vision, taste perversion; Infrequent: tinnitus, amblyopia, deafness; Rare: ophthalmitis, diplopia, photophobia. **Urogenital System**—Frequent: Impotence; Infrequent: metrorrhagia, urinary frequency, urinary tract infection, sexual function abnormal, urinary incontinence, vaginitis; Rare: kidney calculus, abnormal ejaculation, hematuria, menorrhagia, nocturia, polyuria, urinary urgency. **Serious Adverse Events Observed During the Non-US Postmarketing Evaluation of CAMPRAL (acamprosate calcium)** Although no causal relationship to CAMPRAL has been found, the serious adverse event of acute kidney failure has been reported to be temporally associated with CAMPRAL treatment in at least 3 patients and is not described elsewhere in the labeling.

DRUG ABUSE AND DEPENDENCE

Controlled Substance Class Acamprosate calcium is not a controlled substance. Physical and Psychological Dependence CAMPRAL did not produce any evidence of withdrawal symptoms in patients in clinical trials at therapeutic doses. Post marketing data, collected retrospectively outside the U.S., have provided no evidence of CAMPRAL abuse or dependence.

OVERDOSAGE

In all reported cases of acute overdosage with CAMPRAL (total reported doses of up to 56 grams of acamprosate calcium), the only symptom that could be reasonably associated with CAMPRAL was diarrhea. Hypercalcemia has not been reported in cases of acute overdose. A risk of hypercalcemia should be considered in chronic overdosage only. Treatment of overdose should be symptomatic and supportive

Manufactured by: Merck Santé s.a.s. Subsidiary of Merck KGaA, Darmstadt, Germany 37, rue Saint-Romain 69008 LYON FRANCE

Manufactured for FOREST PHARMACEUTICALS, Inc. Subsidiary of Forest Laboratories, Inc. St. Louis, MO 63045 07/04



Legislators Swarm CSAM's Leg Day IV

Donald J. Kurth, M.D., FASAM

alifornia legislators have found the value of speaking to addiction doctors and proved it by making a strong showing at the Fourth Annual Legislative Day sponsored by the California Society of Addiction Medicine. All told, CSAM members met with 32 Assembly members or their aides, as well as representatives of the Governor's office. "The exciting part," explained CSAM President David Pating, "is that the legislators are beginning to compete for our attention. Each one wants to outdo colleagues."

This year's Leg Day was co-sponsored by the California Association of Alcohol and Drug Abuse Counselors. CSAM members welcomed the counselors' involvement in shaping the public policies that affect all of us.

The day began with a meeting of legislative advocates and representatives of CSAM's public policy allies, led by CSAM Public Policy Chair Denise Greene, M.D. Following the opening remarks, ASAM President Elizabeth Howell, M.D., FASAM, addressed the group of 70 attendees and thanked everyone for the inspiration California has provided for all ASAM members in the area of legislative advocacy and public policy. Peter Banys, M.D., then spoke about Proposition 36; Christy Waters, M.D., and Timmen Cermak, M.D., discussed methamphetamine; CSAM Immediate Past President Donald J. Kurth, M.D., FASAM, addressed the human rights of patients with addictive disorders; Diane Sylvestre, M.D., discussed hepatitis C; Gary Jaeger, M.D., FASAM, provided an update on parity; and Warren Daniels, CAADAC, described addiction counselor certification.

After a quick box lunch, the group divided into teams and headed for the Capitol. In addition to the four legislators who addressed the group in the morning, CSAM members met with 28 legislators and/or their aides.

Some legislative districts are affected more than others by the spread of addictive disease and the acute need for increased treatment services. For example, the Fresno area, represented by Assembly Member Bill Maze, has the highest density of intravenous drug use in the Nation. The good news is that Assembly Member Maze is willing to roll up his sleeves and tackle the problem.

Assembly member Jackie Goldberg has


fought for humane detoxification in correctional institutions for years and has carried bills for CSAM in the past. Assembly member Gloria Negrete McCleod is a long-time friend of CSAM and has worked hard to ensure adequate testing and treatment of hepatitis C in incarcerated persons. Assembly member Mark Leno has been a strong advocate for equal protection under the law for all Californians.

The take-home message is that CSAM is making a difference! We are having an impact on public policy and legislation in the State of California and we want you to join

us. Come with us to Sacramento for CSAM Legislative Day V. Watch the CSAM website (www.CSAM-ASAM.org) for details. Better yet, start a Legislative Day in your own home state and help us make a difference for all Americans suffering from addictive disease. Together, we can change the world!

DR. KURTH is Immediate Past President of the California Society of Addiction Medicine and currently serves as Treasurer of the American Society of Addiction Medicine. He also Chairs ASAM's Legislative Advocacy Committee.

TENNESSEE SOCIETY REACTIVATED



Roland Gray, M.D. (center), Director of the Tennessee Medical Foundation physician health program, recently accepted a \$5,000 donation from the Tennessee Society of Addiction Medicine. Making the presentation are TNSAM President Richard G. Soper, M.D., J.D. (at right), and Secretary Jon Butler, M.D. The donation was the first action taken by the Tennessee Society at the chapter's reactivation meeting in December 2005.

STATE SOCIETY & CHAPTER NEWS

Chapter Development Tied to Specialty Recognition

Kevin Kunz, M.D., FASAM

SAM's mission includes the promotion of Ahigh quality scientific and compassionate care to our patients, their families, and our communities. Another essential, yet unfulfilled element of our officially stated mission has been to ensure that our specialty takes a formal seat at the table of mainstream medicine — by gaining recognition from the American Board of Medical Specialties. Addiction medicine needs a presence in every community and state, and as we strive towards national recognition and inclusion the participation of a vibrant and well-organized network of local addiction physicians is necessary. ASAM has two arms, a strong national leadership and local involvement at he state level. The latter must become fully integrated and administratively mature within ASAM if we are to fulfill our mission.

To achieve this, ASAM must transform its state chapters into functioning state medical specialty societies that operate on an equal footing with the state affiliates of other medical specialty groups. This has been the primary goal of the State Medical Specialty Society (SMSS) Program, organized in 2002 under the chairmanship of Richard Beach, M.D. The objectives of the SMSS program are: (1) to create an administrative office within each participating state or ASAM Region, (2) to obtain that state's medical society's formal recognition as a medical specialty society in addiction medicine, and (3) to offer an annual continuing medical education conferences.

We have learned from this project that to achieve these objectives, a state society must take action in six areas: corporate identity, fiscal resources, administration, member activity, communication, and medical society inclusion. The first three are the infrastructure, while the next three are how the members become engaged in carrying out the mission. These are start-up tasks, which many physicians find onerous, but with assistance and guidance, as has been provided via the SMSS program, are achievable. The model each chapter seeks to emulate is that of the California Society of Addiction Medicine (CSAM), the first and most active of ASAM's chapters. CSAM has 350 members and an elected Executive Council, is administered by an association management firm, offers an annual CME conference attended by more than 300, publishes a bi-monthly newsletter and an annual directory, is active in the California Medical Association, and provides education and consultation to the health care community and policymakers. While few states have the capacity to replicate the scale of CSAM's successes, the SMSS pilot program has proven that any state, even those with just a small team of ASAM physicians, can organize successfully and positively impact addictions care at every level, from the bedside to the legislature. Our experience indicates that a quiescent chapter, with a few dedicated leaders, can complete the basic objectives in as little as 18 months.

ACHIEVEMENTS TO DATE

Under the leadership of former ASAM Executive Vice President and current program consultant James F. Callahan, DPA., an impressive list of successes have already been achieved. ASAM physicians are promoting treatment for their patients just as internists, psychiatrists and other specialists form specialty societies to represent their patients' needs within the state. Twenty-seven states now take part in this program, up from 23 in 2004, and 12 in 2002. Our goal is to increase this number to 32 participating states by December 2006, when the current



Members of the SMSS program and the Chapters Council met with ASAM's elected leadership January 20-21st in Bethesda, Maryland. Pictured are:

Front Row, left to right: Michael M. Miller, M.D. (ASAM President-Elect), Richard Soper, M.D. (Tennessee), Elizabeth F. Howell, M.D. (ASAM President), Martha Wunsch, M.D. (Region V Director), Merrill Herman, M.D. (New York), Berton Toews, M.D. (Northwest Chapter), and J ohn Femino, M.D. (Region III Director).

Second Row, left to right: Helen Huff and Nancy Brighindi (ASAM staff), Robert Jones, M.D. (Texas), J. Ramsay Farah, M.D. (Maryland), Donald J. Kurth, M.D. (California and ASAM Treasurer), George D. Hall, M.D. (North Carolina), and Ronald J. Schwerzler, M.D. (Oregon).

Third Row, left to right: Allan M. Ebert, M.D. (Michigan), Valerie Foote (ASAM staff), Kevin Kunz, M.D. (Hawaii and Council Chair), Eileen McGrath, J.D. (ASAM EVP), Gary D. Helmbrecht, M.D. (Virginia), Tim Fischer, D.O. (South Carolina), James F. Callahan, Ph.D. (SMSS Consultant), and C. Chapman Sledge, M.D. (Region X Director).

Back Row, left to right: Wayne E. Pasanen, M.D. (Massachusetts), James M. Miner, M.D. (Virginia), Thomas L. Haynes, M.D. (Region VI Director), George L. Carlson, M.D. (Hawaii), K. L. Hallman, M.D. (Virginia), and Stephen A. Wyatt, D.O. (Connecticut).

Not pictured are Jeffrey C. Craddock, M.D. (South Carolina), Miriam H. Komaromy, M.D. (New Mexico), Sarz Maxwell, M.D. (Illinois), David Pating, M.D. (California), John D. Patz, M.D. (Washington State), Mark Publicker, M.D. (Maine), Marvin Seppala, M.D. (Region VIII Director) and Howard Wetsman, M.D. (Region VII Director).

STATE SOCIETY & CHAPTER NEWS

funding ends, and a new budgeting strategy will be needed.

Most patients with addictive disorders are unlikely to receive adequate medical care until addiction medicine is part of our country's health and medical education systems. The ultimate goal for assuring this is to have an American Board of Addiction Medicine as a member board of the American Board of Medical Specialties. Formal recognition of addiction medicine by the AMA and the state medical societies is an essential step toward this goal. Through the efforts of the SMSS and its state affiliates, 10 state medical societies now recognize their state society of addiction medicine and have given addiction medicine a seat in the state society's house of delegates or governing body. This is up from 7 in 2004, and 3 in 2003. Our goal for 2006 is to increase the number to 15. We have demonstrated that addiction medicine can be a welcome and functional member of the family of medical specialties.

In participating states, the number of medical education conferences has increased significantly each year. In Texas, attendance at the Region VII 2005 annual conference tripled over 2004, despite the difficulties posed by hurricane Katrina. In 2006, a 1.5 day physician education track will be added to the Cape Cod Symposium (Massachusetts), an annual conference that draws 500 registrants. Several states are routinely incorporating addiction medicine sessions with their state societies annual meeting.

Public policy achievements have been equally remarkable, including successes in battling against state bills to legalize marijuana (under the euphemism of "medical marijuana") and in working for the repeal of harmful legislation such as the Uniform Accident and Sickness Policy Provision Law [UPPL]. The UPPL is a major barrier to screening and referral for alcohol and other drug problems because it expressly states that "the (health) insurer shall not be liable for any loss sustained or contracted in consequence of the insured's being intoxicated or under the influence of any narcotic, unless administered on the advice of a physician." Parity for addiction medicine has also been addressed at the state level.

GOALS FOR THE FUTURE

Among our goals for 2006, two are essential to assure the continuation of the program. The first is to help each state society secure sufficient income and administrative staff to support medical education and public policy. The second goal is to integrate the SMSS program into the ASAM operating budget under the ASAM Chapters Council, which is charged with chapter development and growth. It is clear that the Chapters Council must continue to encourage and support the objectives of the SMSS program for the current momentum to continue.

Robert L. DuPont, M.D., FASAM, continues to offer his wisdom and has kept us focused on what we are trying to achieve. As he does so well, he summarized our purpose when he said, "Our work...is not for the glory of ASAM. Our work is aimed at helping the thousands of addicted people, in communities large and small all over the country, find their ways out of addiction into recovery through the use of modern, scientific addiction medicine. ASAM member physicians in those communities are the ones who can best help their patients find their ways to recovery. They are the dedicated physicians who are best able to give their patients the advantages of the evidence-based medicine that is newly available to facilitate lasting recovery, including wholehearted support of lifelong participation in Alcoholics Anonymous and Narcotics Anonymous as the best way to prevent relapses into the quicksand of addiction."

In summary, the strength of ASAM is found in our two arms: a strong national leadership and involvement at the state level. Both are required to fulfill our mission. Indeed, it is the hands-on efforts of our members in their hometowns and home states that ultimately increase our patients' access to high quality care. It is at the local level where we directly interface with, assist and educate our peers and the public. If addiction medicine is to reach full specialty status, the input and enthusiasm of a diverse and included membership will be essential.

ASAM is grateful to John P. McGovern, M.D., FASAM (Honorary) and the John P. McGovern Foundation, as well as to the Scaife Family Foundation, for their steadfast support of this vital initiative. In addition, the ASAM Board, which initially approved the program in 2002, has been unwavering in its support of the initiative, as has Executive Vice President/CEO Eileen McGrath, J.D.

Dr. Kunz chairs the SMSS program and the ASAM Chapters Council. He is Past-President of the Hawaii Society of Addiction Medicine.

CTSAM Works for Better Care of Physicians

Stephen A. Wyatt, D.O.

Douglas Gibson, M.D., Medical Director of the Connecticut Physician Health Program, has made tremendous strides in establishing a more science-based, equitable, and confidential program to assist impaired health care professionals in Connecticut. The vehicle is Raised Bill No. 5718 [LCO 2532], An Act Concerning a Professional Assistance Program for Health Care Professionals. The language of the bill can be accessed online at HTTP://WWW.CGA.CT.GOV/ 2006/TOB/H/2006HB-05718-R00-HB.HTM

This important measure currently is before the Committee on Public Health of the state legislature. Please review the bill, discuss it with your colleagues, and contact your legislators and friends to advocate for its enactment.

Dr. Gibson's work is an example of the progress we are making in Connecticut. The Connecticut Society of Addiction Medicine is at the table in the state medical society. CTSAM is the organization the public and the legislature turn to most frequently for expertise in our field. And we have very active representation nationally through ASAM — the most active voice in the field of addiction medicine. There has never been a time when addiction medicine has had such an opportunity to lead; indeed, with the advent of new therapeutic options and the growth of Fellowship training, physicians are now the lynchpins of the therapeutic team.


I'm also pleased to announce that our state society has a new website, www. CTSAM.ORG. Please consider the importance of supporting this strong and active voice by joining or renewing your membership in the Connecticut Society and ASAM. Together we can fight for appropriate care for our patients and educate the public and policymakers. Treat addiction, save lives!

DR. WYATT is President of the Connecticut Society of Addiction Medicine and Medical Director of the Stonington Institute, North Stonington, CT.

Treat the Condition

Opioid Dependence Is a Chronic Medical Condition

Long-term, fundamental changes to structure and function of the brain occur.^{1,2}



Intravenous misuse of buprenorphine, usually in combination with benzodiazepines or other CNS depressants, has been associated with significant respiratory depression and death.

SUBOXONE has potential for abuse and produces dependence of the opioid type with a milder withdrawal syndrome than full agonists.

Cytolytic hepatitis and hepatitis with jaundice have been observed in the addicted population receiving buprenorphine.

There are no adequate and well-controlled studies of SUBOXONE (a category C medication) in pregnancy.

Due caution should be exercised when driving cars or operating machinery.

The most commonly reported adverse events with SUBOXONE include: headache (36%, placebo 22%), withdrawal syndrome (25%, placebo 37%), pain (22%, placebo 19%), nausea (15%, placebo 11%), insomnia (14%, placebo 16%), and sweating (14%, placebo 10%).

Please see adjacent Brief Summary of Prescribing Information.

References: 1. Leshner Al, Koob GF. Drugs of abuse and the brain. Proc Assoc Am Physicians. 1999;111(2):99-108. 2. Leshner Al, Addiction is a brain disease, and it matters. Science. 1997;278:45-47.

Transform the Life



In the Privacy and Convenience of Your Office

SUBOXONE, combined with counseling, can be used to treat opioid-dependent patients with privacy,* as other chronic, medical conditions are treated.

Target the Biological Basis of Opioid Dependence

SUBOXONE suppresses withdrawal symptoms, decreases cravings, and improves treatment retention. With the support of pharmacotherapy and counseling, patients may gain control over opioid dependence and be able to address other aspects of their lives.

To learn more, call 1-877-SUBOXONE or visit suboxone.com

*Under the Drug Addiction and Treatment Act of 2000 (DATA 2000), physicians who meet certain qualifying requirements may prescribe SUBOXONE. **Visit OpioidDependence.com** for information about qualifying.



SUBOXONE (CIII)

(buprenorphine HCl and naloxone HCl dihydrate sublingual tablets)

SUBUTEX (CIII)

(buprenorphine HCI sublingual tablets)

Ry only

Brief Summary: Consult the SUBOXONE package insert for complete prescribing information.

Under the Drug Addiction Treatment Act of 2000 (DATA) codified at 21 U.S.C. 823(g), prescription use of this product in the treatment of opioid dependence is limited to physicians who meet certain qualifying requirements, and have notified the Secretary of Health and Human Services (HHS) of their intent to prescribe this product for the treatment of onioid dependence.

INDICATIONS AND USAGE

SUBOXONE and SUBUTEX are indicated for the treatment of opioid dependence

CONTRAINDICATIONS

SUBOXONE and SUBUTEX should not be administered to patients who have been shown to be hypersensitive to buprenorphine, and SUBOXONE should not be administered to patients who have been shown to be hypersensitive to naloxone.

WARNINGS

Respiratory Depression: Significant respiratory depression has been associated with buprenorphine, particularly by the intravenous route. A number of deaths have occurred when addicts have intravenously misused buprenorphine usually with berzodiazepines concomitantly. Deaths have also been reported in association with concomitant administration of buprenorphine with other depressants such as alcohol or other opioids. Patients should be warned of the potential danger of the self-administration of benzodiazepines or other depressants while under treatment with SUBUTEX or SUBOXONE.

IN THE CASE OF OVERDOSE, THE PRIMARY MANAGEMENT SHOULD BE THE RE-ESTABLISHMENT OF ADEQUATE VENTILATION WITH MECHANICAL ASSISTANCE OF RESPIRATION, IF REQUIRED. NALOXONE MAY NOT BE EFFECTIVE IN REVERSING ANY RESPIRATORY DEPRESSION PRODUCED BY BUPRENORPHINE.

SUBOXONE and SUBUTEX should be used with caution in patients with compromised respiratory function (e.g., chronic obstructive pulmonary disease, cor pulmonale, decreased respiratory reserve, hypoxia, hypercapnia, or pre-existing respiratory decression.

CNS Depression: Patients receiving buprenorphine in the presence of other narcotic analgesics, general anesthetics, benzodiazepines, phenothiazines, other tranquilizers, sedative/hypnotics or other CNS depressants (including alcohol) may exhibit increased CNS depression. When such combined therapy is contemplated, reduction of the dose of one or both agents should be considered.

Drug Abuse and Dependence: SUBOXONE and SUBUTEX are controlled as Schedule III narcotics under the Controlled Substances Act.

Buprenorphine is a partial agonist at the mu-opioid receptor and chronic administration produces dependence of the opioid type, characterized by moderate withdrawal upon abrupt discontinuation or rapid taper. The withdrawal syndrome is milder than seen with full agonists, and may be delayed in onset.

Neonatal withdrawal has been reported in the infants of women treated with SUBUTEX during pregnancy (See PRECALITIONS)

SUBOXONE contains naloxone and if misused parenterally, is highly likely to produce marked and intense withdrawal symptoms in subjects dependent on other opioid agonists.

Hepatitis, Hepatic Events: Cases of cytolytic hepatitis and hepatitis with jaundice have been observed in the addict population receiving buprenorphine both in clinical trials and in post-marketing adverse event reports. The spectrum of abnormalities ranges from transient asymptomatic elevations in hepatic transaminases to case reports of hepatic failure, hepatic necrosis, hepatorenal syndrome, and hepatic encephalopathy. In many cases, the presence of pre-existing liver enzyme abnormalities, infection with hepatitis B or hepatitis C virus, concomitant usage of other potentially hepatotoxic drugs, and ongoing injecting drug use may have played a causative or contributory role. In other cases, insufficient data were available to determine the etiology of the abnormality. The possibility exists that buprenorphine had a causative or contributory role in the development of the hepatic abnormality in some cases. Measurements of liver function tests during treatment is also recommended. A biological and etiological evaluation is recommended when a hepatic event is suspected. Depending on the case, the drug should be carefully discontinued to prevent withdrawal symptoms and a return to illicit drug use, and strict monitoring of the patient should be initiated.

Allergic Reactions: Cases of acute and chronic hypersensitivity to buprenorphine have been reported both in clinical trials and in the post-marketing experience. The most common signs and symptoms include rashes, hives, and pruntus. Cases of bronchospasm, angioneurotic edema, and anaphylactic shock have been reported. A history of hypersensitivity to buprenorphine is a contraindication to SUBUTEX or SUBOXONE use. A history of hypersensitivity to naloxone is a contraindication to SUBOXONE use.

Use in Ambulatory Patients: SUBOXONE and SUBUTEX may impair the mental or physical abilities required for the performance of potentially dangerous tasks such as driving a car or operating machinery, especially during drug induction and dose adjustment. Patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that buprenorphine therapy does not adversely affect their ability to engage in such activities. Like other opioids, SUBOXONE and SUBUTEX may produce orthostatic hypotension in ambulatory patients.

Head Injury and Increased Intracranial Pressure: SUBOXONE and SUBUTEX, like other potent opioids, may elevate cerebrospinal fluid pressure and should be used with caution in patients with head injury, intracranial lesions and other circumstances where cerebrospinal pressure may be increased. SUBOXONE and SUBUTEX can produce miosis and changes in the level of consciousness that may interfere with patient evaluation.

Opioid Withdrawal Effects: Because it contains naloxone, SUBOXONE is highly likely to produce marked and intense withdrawal symptoms if misused parenterally by individuals dependent on opioid agonists such as heroin, morphine, or methadone. Sublingually, SUBOXONE may cause opioid withdrawal symptoms in such persons if administered before the agonist effects of the opioid have subsided.

PRECAUTIONS

General: SUBOXONE and SUBUTEX should be administered with caution in elderly or debilitated patients and those with severe impairment of hepatic, pulmonary, or renal function; myxedema or hypothyroidism, adrenal cortical insufficiency (e.g., Addison's disease); CNS depression or coma; toxic psychoses; prostatic hypertrophy or urethral stricture; acute alcoholism; delirium tremens; or kyphoscoliosis.

The effect of hepatic impairment on the pharmacokinetics of buprenorphine and naloxone is unknown. Since both drugs are extensively metabolized, the plasma levels will be expected to be higher in patients with moderate and severe hepatic impairment. However, it is not known whether both drugs are affected to the same degree. Therefore, dosage should be adjusted and patients should be watched for symptoms of precipitated opioid withdrawal.

Buprenorphine has been shown to increase intracholedochal pressure, as do other opioids, and thus should be administered with caution to patients with dysfunction of the biliary tract.

As with other mu-opioid receptor agonists, the administration of SUBOXONE or SUBUTEX may obscure the diagnosis or clinical course of patients with acute abdominal conditions.

Drug Interactions: Buprenorphine is metabolized to norbuprenorphine by cytochrome CYP 3A4. Because CYP 3A4 inhibitors may increase plasma concentrations of buprenorphine, patients already on CYP 3A4 inhibitors such as azole antifungals (e.g., ketoconazole), macrolide antibiotics (e.g., erythromycin), and HIV protease inhibitors (e.g., ritonavir, indinavir and saquinavir) should have their dose of SUBUTEX or SUBOXONE adjusted.

Based on anecdotal reports, there may be an interaction between buprenorphine and benzodiazepines. There have been a number of reports in the post-marketing experience of coma and death associated with the concomitant intravenous misuse of buprenorphine and benzodiazepines by addicts. In many of these cases, buprenorphine was misused by self-injection of crushed SUBUTEX tablets. SUBUTEX and SUBOXONE should be prescribed with caution to patients on benzodiazepines or other drugs that act on the central nervous system, regardless of whether these drugs are taken on the advice of a physician or are taken as drugs of abuse. Patients should be warned of the potential danger of the intravenous self-administration of benzodiazepines while under treatment with SUBOXONE or SUBUTEX.

Information for Patients: Patients should inform their family members that, in the event of emergency, the treating physician or emergency room staff should be informed that the patient is physically dependent on narcotics and that the patient is being treated with SUBOXONE or SUBUTEX.

Patients should be cautioned that a serious overdose and death may occur if benzodiazepines, sedatives, tranquilizers, antidepressants, or alcohol are taken at the same time as SUBOXONE or SUBUTEX.

SUBOXONE and SUBUTEX may impair the mental or physical abilities required for the performance of potentially dangerous tasks such as driving a car or operating machinery, especially during drug induction and dose adjustment. Patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that buprenorphine therapy does not adversely affect their ability to engage in such activities. Like other oploids, SUBOXONE and SUBUTEX may produce orthostatic hypotension in ambulatory patients.

Patients should consult their physician if other prescription medications are currently being used or are prescribed for future use

Carcinogenesis, Mutagenesis and Impairment of Fertility: Carcinogenicity. Carcinogenicity data on SUBOXONE are not available. Carcinogenicity studies of buprenorphine were conducted in Sprague-Dawley rats and CD-1 mice Buprenorphine was administered in the diet to rats at doses of 0.6, 5.5, and 5.6 m/kg/day (estimated exposure was approximately 0.4, 3 and 35 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis) for 27 months. Statistically significant dose-related increases in testicular interstitial (Leydig's) cell tumors occurred according to the trend test adjusted for survival. Pair-wise comparison of the high dose against control failed to show statistical significance. In an 86-week study in CD-1 mice, buprenorphine was not carcinogenic at dietary doses up to 100 mg/kg/day (estimated exposure was approximately 30 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis).

Mutagenicity: SUBOXONE: The 4:1 combination of buprenorphine and naloxone was not mutagenic in a bacterial mutation assay (Ames test) using four strains of *S. typhimurium* and two strains of *E. coli*. The combination was not clastogenic in an *in vitro* cytogenetic assay in human lymphocytes, or in an intravenous micronucleus test in the rat. SUBUTEX: Buprenorphine was studied in a series of tests utilizing gene, chromosome, and DNA interactions in both prokaryotic and eukaryotic systems. Results were negative in yeast (*Saccharomyces cerevisiae*) for recombinant, gene convertant, or forward mutations; negative in *Bacillus subtilis* "rec" assay, negative for clastogenicity in CHOells, Chinese hamster bone marrow and spermatogonia cells, and negative in the mouse lymphoma L5178Y assay. Results were equivocal in the Ames test: negative in studies in two laboratories, but positive for frame shift mutation at a high dose (5 mg/plate) in a third study. Results were positive in the Green-Tweets (*E. coli*) survival test, positive in a DNA synthesis inhibition (DSI) test with testicular tissue from mice, for both *in vivo* and *in vitro* incorporation of [H]thymidine, and positive in unscheduled DNA synthesis (UDS) test using testicular cells from mice.

Impairment of Fertility: SUBOXONE: Dietary administration of SUBOXONE in the rat at dose levels of 500 ppm or greater (equivalent to approximately 47 mg/kg/day or greater; estimated exposure was approximately 28 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis) produced a reduction in fertility demonstrated by reduced female conception rates. A dietary dose of 100 ppm (equivalent to approximately 10 mg/kg/day; estimated exposure was approximately 6 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis) had no adverse effect on fertility.

SUBUTEX: Reproduction studies of buprenorphine in rats demonstrated no evidence of impaired fertility at daily oral doses up to 80 mg/kg/day (estimated exposure was approximately 50 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis) or up to 5 mg/kg/day *im* or *sc* (estimated exposure was approximately 3 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis).

Pregnancy: Pregnancy Category C:

Teratogenic effects: SUBOXONE: Effects on embryo-fetal development were studied in Sprague-Dawley rats and Russian white rabbits following oral aft (1:1) and intramuscular (3:2) administration of mixtures of buprenorphine and aloxone. Following oral administration to rats and rabbits, no teratogenic effects were observed at doses up to 250 mg/kg/day and 40 mg/kg/day, respectively (estimated exposure was approximately 150 times and 50 times, respectively, the recommended human daily sublingual dose of 16 mg on a mg/m² basis). No definitive drug-related teratogene effects were observed in rats and rabbits at intramuscular doses up to 30 mg/kg/day (estimated exposure was approximately 20 times and 35 times, respectively, the recommended human daily dose of 16 mg on a mg/m² basis). Acephalus was observed in one rabbit fetus from the low-dose group and omphacele was observed in the variet fetuses from the same litter in the mid-dose group; no findings were observed in fetuses from the high-dose group. Following oral administration to the rat, dose-related post-implantation losses, evidenced by increases in the numbers of early resorptions with consequent reductions in the numbers of fetuses, were observed at doses of 10 mg/kg/day or greater (estimated exposure was approximately 6 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis). In the rabbit, increased post-implantation losses occurred at an oral dose of 40 mg/kg/day. Following intramuscular administration in the rat and the rabbit, post-implantation losses, as evidenced by decreases in live fetuses and increases in resorptions, occurred at 30 mg/kg/day.

SUBUTEX: Buprenorphine was not teratogenic in rats or rabbits after $\stackrel{'}{m}$ or sc doses up to 5 mg/kg/day (estimated exposure was approximately 3 and 6 times, respectively, the recommended human daily sublingual dose of 16 mg on a mg/m² doses up to 0.8 mg/kg/day (estimated exposure was approximately 0.5 times and equal to, respectively, the recommended human daily sublingual dose of 16 mg on a mg/m² basis), or after oral doses up to 160 mg/kg/day in rats (estimated exposure was approximately 9.5 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis) and 25 mg/kg/day in rabbits (estimated exposure was approximately 30 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis). Significant increases in skeletal abnormalities (e.g., extar thoracic vertebra or thoraco-lumbar ribs) were noted in rats after sc administration of 1 mg/kg/day and up (estimated exposure was approximately 0.6 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis), but were not observed at oral doses up to 160 mg/kg/day, increases in skeletal abnormalities in rabbit after sc administration of 5 mg/kg/day (scimated exposure was approximately 6.6 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis) or oral administration of 1 mg/kg/day or greater (estimated exposure was approximately 6.9 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis) were not statistically significant.

In rabbits, buprenorphine produced statistically significant pre-implantation losses at oral doses of 1 mg/kg/day or greater and post-implantation losses that were statistically significant at in doses of 0.2 mg/kg/day or greater (estimated exposure was approximately 0.3 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis). There are no adequate and well-controlled studies of SUBOXONE or SUBUTEX in pregnant women. SUBOXONE or SUBUTEX should only be used during pregnancy if the potential benefit justifies the potential risk to the fetus.

Non-teratogenic effects: Dystocia was noted in pregnant rats treated im with buprenorphine 5 mg/kg/day (approximately 3 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis). Both fertility and peri- and postnatal development studies with buprenorphine in rats indicated increases in neonatal mortality after oral doses of 0.8 mg/kg/day and up (approximately 0.5 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis), after im doses of 0.5 mg/kg/day and up (approximately 0.3 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis), and after sc doses of 0.1 mg/kg/day and up (approximately 0.06 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis). Delays in the occurrence of righting reflex and startle response were noted in rat pups at an oral dose of 80 mg/kg/day (approximately 50 times the recommended human daily sublingual dose of 16 mg on a mg/m² basis).

Neonatal Withdrawal: Neonatal withdrawal has been reported in the infants of women treated with SUBUTEX during pregnancy. From post-marketing reports, the time to onset of neonatal withdrawal symptoms ranged from Day 1 to Day 8 of life with most occurring on Day 1. Adverse events associated with neonatal withdrawal syndrome included hypertonia, neonatal remor, neonatal agitation, and myoclonus. There have been rare reports of convulsions and in one case, apnea and bradycardia were also reported.

Nursing Mothers: An apparent lack of milk production during general reproduction studies with buprenorphine in rats caused decreased viability and lactation indices. Use of high doses of sublingual buprenorphine in pregnant women showed that buprenorphine passes into the mother's milk. Breast-feeding is therefore not advised in mothers treated with SUBUTEX or SUBOXONE.

Pediatric Use: SUBOXONE and SUBUTEX are not recommended for use in pediatric patients. The safety and effectiveness of SUBOXONE and SUBUTEX in patients below the age of 16 have not been established.

ADVERSE REACTIONS

The safety of SUBOXONE has been evaluated in 497 opioid-dependent subjects. The prospective evaluation of SUBOXONE was supported by clinical trials using SUBUTEX (buprenorphine tablets without naloxone) and other trials using buprenorphine sublingual solutions. In total, safety data are available from 3214 opioid-dependent subjects exposed to buprenorphine at doses in the range used in treatment of opioid addiction.

Few differences in adverse event profile were noted between SUBOXONE and SUBUTEX or buprenorphine administered as a sublingual solution.

In a comparative study, adverse event profiles were similar for subjects treated with 16 mg SUBOXONE or 16 mg $\,$ SUBUTEX. The following adverse events were reported to occur by at least 5% of patients in a 4-week study (Table 1).

Table 1. Adverse Events (≥5%) by Body System and Treatment Group in a 4-week Study

·	N (%)	N (%)	N (%)
Body System /Adverse Event (COSTART Terminology)	SUBOXONE 16 mg/day N=107	SUBUTEX 16 mg/day N=103	Placebo N=107
Body as a Whole			
Asthenia	7 (6.5%)	5 (4.9%)	7 (6.5%)
Chills	8 (7.5%)	8 (7.8%)	8 (7.5%)
Headache	39 (36.4%)	30 (29.1%)	24 (22.4%)
Infection	6 (5.6%)	12 (11.7%)	7 (6.5%)
Pain	24 (22.4%)	19 (18.4%)	20 (18.7%)
Pain Abdomen	12 (11.2%)	12 (11.7%)	7 (6.5%)
Pain Back	4 (3.7%)	8 (7.8%)	12 (11.2%)
Withdrawal Syndrome	27 (25.2%)	19 (18.4%)	40 (37.4%)
Cardiovascular System			
Vasodilation	10 (9.3%)	4 (3.9%)	7 (6.5%)
Digestive System			
Constipation	13 (12.1%)	8 (7.8%)	3 (2.8%)
Diarrhea	4 (3.7%)	5 (4.9%)	16 (15.0%)
Nausea	16 (15.0%)	14 (13.6%)	12 (11.2%)
Vomiting	8 (7.5%)	8 (7.8%)	5 (4.7%)
Nervous System			
Insomnia	15 (14.0%)	22 (21.4%)	17 (15.9%)
Respiratory System			
Rhinitis	5 (4.7%)	10 (9.7%)	14 (13.1%)
Skin and Appendages			
Sweating	15 (14.0%)	13 (12.6%)	11 (10.3%)

The adverse event profile of buprenorphine was also characterized in the dose-controlled study of buprenorphine solution, over a range of doses in four months of treatment. Table 2 shows adverse events reported by at least 5% of subjects in any dose group in the dose-controlled study.

Table 2. Adverse Events (≥5%) by Body System and Treatment Group in a 16-week Study

	Buprenorphine Dose*				
Body System/Adverse	Very Low* Low* Moderate* H		High*	Total*	
Event (COSTART	(N=184)	(N=180)	(N=186)	(N=181)	(N=731)
Terminology)	N (%)	N (%)	N (%)	N (%)	N (%)
Body as a Whole					
Abscess	9 (5%)	2 (1%)	3 (2%)	2 (1%)	16 (2%)
Asthenia	26 (14%)	28 (16%)	26 (14%)	24 (13%)	104 (14%)
Chills	11 (6%)	12 (7%)	9 (5%)	10 (6%)	42 (6%)
Fever	7 (4%)	2 (1%)	2 (1%)	10 (6%)	21 (3%)
Flu Syndrome	4 (2%)	13 (7%)	19 (10%)	8 (4%)	44 (6%)
Headache	51 (28%)	62 (34%)	54 (29%)	53 (29%)	220 (30%)
Infection	32 (17%)	39 (22%)	38 (20%)	40 (22%)	149 (20%)
Injury Accidental	5 (3%)	10 (6%)	5 (3%)	5 (3%)	25 (3%)
Pain	47 (26%)	37 (21%)	49 (26%)	44 (24%)	177 (24%)
Pain Back	18 (10%)	29 (16%)	28 (15%)	27 (15%)	102 (14%)
Withdrawal Syndrome	45 (24%)	40 (22%)	41 (22%)	36 (20%)	162 (22%)
Digestive System					
Constipation	10 (5%)	23 (13%)	23 (12%)	26 (14%)	82 (11%)
Diarrhea	19 (10%)	8 (4%)	9 (5%)	4 (2%)	40 (5%)
Dyspepsia	6 (3%)	10 (6%)	4 (2%)	4 (2%)	24 (3%)
Nausea	12 (7%)	22 (12%)	23 (12%)	18 (10%)	75 (10%)
Vomiting	8 (4%)	6 (3%)	10 (5%)	14 (8%)	38 (5%)
Nervous System					
Anxiety	22 (12%)	24 (13%)	20 (11%)	25 (14%)	91 (12%)
Depression	24 (13%)	16 (9%)	25 (13%)	18 (10%)	83 (11%)
Dizziness	4 (2%)	9 (5%)	7 (4%)	11 (6%)	31 (4%)
Insomnia	42 (23%)	50 (28%)	43 (23%)	51 (28%)	186 (25%)
Nervousness	12 (7%)	11 (6%)	10 (5%)	13 (7%)	46 (6%)
Somnolence	5 (3%)	13 (7%)	9 (5%)	11 (6%)	38 (5%)
Respiratory System					
Cough Increase	5 (3%)	11 (6%)	6 (3%)	4 (2%)	26 (4%)
Pharyngitis	6 (3%)	7 (4%)	6 (3%)	9 (5%)	28 (4%)
Rhinitis	27 (15%)	16 (9%)	15 (8%)	21 (12%)	79 (11%)
Skin and Appendages					
Sweat	23 (13%)	21 (12%)	20 (11%)	23 (13%)	87 (12%)
Special Senses					
Runny Eyes	13 (7%)	9 (5%)	6 (3%)	6 (3%)	34 (5%)

^{*}Sublingual solution. Doses in this table cannot necessarily be delivered in tablet form, but for comparison numoses:

OVERDOSAGE

Manifestations: Manifestations of acute overdose include pinpoint pupils, sedation, hypotension, respiratory depression and death

Treatment: The respiratory and cardiac status of the patient should be monitored carefully. In the event of depression of respiratory or cardiac function, primary attention should be given to the re-establishment of adequate respiratory exchange through provision of a patent airway and institution of assisted or controlled ventilation. Oxygen, intravenous fluids, vaso-pressors, and other supportive measures should be employed as indicated.

IN THE CASE OF OVERDOSE, THE PRIMARY MANAGEMENT SHOULD BE THE RE-ESTABLISHMENT OF ADEQUATE VENTILATION WITH MECHANICAL ASSISTANCE OF RESPIRATION. IF REQUIRED. NALOXONE MAY NOT BE EFFECTIVE IN REVERSING ANY RESPIRATORY DEPRESSION PRODUCED BY BUPRENORPHINE.

High doses of naloxone hydrochloride, 10-35 mg/70 kg may be of limited value in the management of buprenorphine overdose. Doxapram (a respiratory stimulant) also has been used.

Manufactured by: Reckitt Benckiser Healthcare (UK) Ltd, Hull, UK, HU8 7DS Distributed by: Reckitt Benckiser Pharmaceuticals, Inc., Richmond, VA 23235 #138274BS July 2005

Membership Report

ASAM's Member-Get-A-Member Campaign Continues

ASAM's Member-Get-A-Member Campaign, which began October 1, 2005, continues through April 15th, so there still is time to take part! Share your ASAM experience when you are meeting with colleagues and ask them to consider joining ASAM. Not only will you be doing yourself a favor by strengthening your Society, but your colleagues will thank you for inviting them.

Each new member you recruit moves you closer to receiving one of the following*:

1st prize: One free registration to the 2007 Med-Sci

Conference (value up to \$585)

2nd prize: One free membership renewal for 2007

(value up to \$485)

3rd prize: One free copy of ASAM's textbook, *Principles of*

Addiction Medicine, 3rd Edition (value up to \$175).

As of February 2006, each of the following members has recruited at least one new member:

RECRUITED 3 NEW MEMBERS

Richard G. Soper, M.D.

RECRUITED 2 NEW MEMBERS

Sanjay Chandragiri, M.D.

RECRUITED 1 NEW MEMBER

Mihran N. Ask, M.D. Jeffrey Baxter, M.D.

Louis E. Baxter, Sr., M.D., FASAM E. Coffman, D.O.

> Barry E. Cole, M.D. Jeff Daiter, M.D.

Christina Delos-Reves, M.D.

Allan Ebert, M.D.

J. Ramsay Farah, M.D.

Marc Galanter, M.D., FASAM Samuel Ganz, M.D.

Frederick C. Goggans, M.D. Lloyd J. Gordon III, M.D., FASAM

Susan T. Howard, M.D., FASAM

Ebenezer Kolade, M.D. Vijava Kotha, M.D.

Kevin B. Kunz, M.D., FASAM David R. McDuff, M.D. Anthony Miller, M.D. Michael M. Miller, M.D.,

FASAM Tania Nordli, M.D.

David Pating, M.D.

Mark R. Publicker, M.D.

Nader Sharifi, M.D. Paul Tisher, M.D.

Scott Treworgy, M.D.

Michael Varenbut, M.D.

Lawrence Westreich, M.D. Melanie Willows, M.D.

Jack Woodside, M.D.

William Wooten, M.D.

Add your name to the growing list and increase your chances of receiving a prize by recruiting a new member today! Visit the ASAM website at www.asam.org for details and membership application

[&]quot;Very low" dose (1 mg solution) would be less than a tablet dose of 2 mg "Low" dose (4 mg solution) approximates a 6 mg tablet dose

[&]quot;Moderate" dose (8 mg solution) approximates a 12 mg tablet dose

[&]quot;High" dose (16 mg solution) approximates a 24 mg tablet dose

^{*} In the event of a tie, the winner will be determined by a drawing.

RUTH FOX MEMORIAL ENDOWMENT FUND

Associate Medical Director

THE NORTH CAROLINA PHYSICIANS HEALTH PROGRAM

(www.ncphp.org) seeks a physician in recovery for the Associate Medical Director position. The AMD helps to facilitate treatment of health care professionals with medical conditions that could compromise patient care.

Qualifications include a medical degree from an AMA-accredited school and public speaking skills. Successful completion of AMArecognized medical residency and certification by ASAM or ABPN are highly desirable. Must quality for NC medical licensure. Send CV, bio, two letters of reference, and salary requirements to:

Kim McCallie **NC Physicians Health Program** 220 Horizon Drive, Suite 218 Raleigh, NC 27615 Email: kim@ncphp.org



Dr. Ruth Fox

Dear Colleague:

The 2006 Ruth Fox Donor Reception is scheduled for 6:30 to 8:30 p.m. on Friday evening, May 5th, during ASAM's Medical-Scientific Conference in San Diego. The reception honors the generosity of those who have made donations to the Fund. As in years past, the cost of the reception has been underwritten by a generous gift from ASAM member Joseph E. Dorsey, M.D., and Mrs. Dorsey.

The Reception also provides an opportunity to celebrate the achievements of this year's recipients of the Ruth Fox Scholar-

ships, given to an outstanding group of physicians-in-training. To date, 24 such scholarships have been awarded. Scholarships cover travel, hotel and registration expenses for recipients to attend ASAM's Annual Medical-Scientific Conference and Ruth Fox Course, as well as one year's free membership in ASAM. The four scholarship recipients for 2006 are Kathleen Ang-Lee, M.D. (Seattle, Washington), Katrina Ball, D.O. (Loma Linda, California), Norana Irene Caivano, M.D. (West Hollywood, California), and Mark Hrymoc, M.D. (Harbor UCLA Medical Center, Los Angeles).

The Endowment was established to create a fiscally sound base so as to assure ASAM's continued ability to realize its mission: to provide ongoing leadership in newly emerging areas affecting the field of addiction medicine, to continue its commitment to educating physicians, to increasing access to care and to improving the quality of care. With the professional and financial support of ASAM's members and friends, the Fund will achieve its mission

Invitations to the Ruth Fox Donor Reception are extended only to donors, so if you have not already contributed or pledged to the Endowment Fund, please do so now. For information about making a pledge, contribution, bequest, memorial tribute, or to discuss other types of gifts in confidence, please contact Claire Osman by phone at 1-800/257-6776 or 1-718/275-7766, or email Claire at ASAMCLAIRE@AOL.COM. She welcomes your calls. All contributions to the Endowment Fund are tax-deductible to the full extent allowed by law.

Max A. Schneider, M.D., FASAM Chair, Ruth Fox Memorial Endowment Subcommittee

Claire Osman Director of Development

ASAM STAFF & CONSULTANTS

Eileen McGrath, J.D. **Executive Vice President/CEO** EMCGRATH@ASAM.ORG

Berit Boegli Conferences & **Meetings Assistant** BBOEG@ASAM.ORG

Nancy Brighindi Director of Membership & Chapter Development NBRIG@ASAM.ORG

Ruby Bailey Edmondson Office Manager/Receptionist RBAIL@ASAM.ORG

Valerie Foote **Data Entry Operator** VFOOT@ASAM.ORG

Joanne Gartenmann Consultant JGART@ASAM.ORG

Except where indicated, all staff can be reached at ASAM's Headquarters Office, 4601 North Park Ave., Suite 101 Upper, Chevy Chase, MD 20814; phone 301/656-3920; EMAIL@ASAM.ORG.

Tracy Gartenmann Director of PCSS & **Buprenorphine Training** TGART@ASAM.ORG

Alexis Geier-Horan Director, Government Relations **Relations Assistant** AGEIER@ASAM.ORG

Gionne Graetz **Bureprenorphine Training** and PCSS Manager AGEIER@ASAM.ORG

Maria Glanz Exec. Assistant to the EVP MGLANZ@ASAM.ORG

Gionne Graetz

Buprenorphine Training & PCSS Manager GGRAETZ@ASAM.ORG

Amy Hotaling Member & Chapter **Development Manager** AHOTA@ASAM.ORG

Lynda Jones Director of Finance LIONE@ASAM.ORG

Sandra Metcalfe Acting Director, **Conferences & Meetings** SMETC@ASAM.ORG

Claire Osman

Director of Development Phone: 1-800/257-6776 Fax: 718/275-7666 ASAMCLAIRE@AOL.COM

Noushin Shariate Accounts Payable NSHAR@ASAM.ORG

Christopher Weirs Director of Credentialing/ IT Manager CWEIR@ASAM.ORG

Bonnie B. Wilford Managing Editor, **ASAM Publications** 29261 Pin Oak Way Easton, MD 21601-4631 Phone: 410/770-4866 Fax: 410/770-4711 BBWILFORD@AOL.COM




Today's source for tomorrow/s pain management!

ESP Tool Kit

A multimedia collection of practical, educational resources and tools, focusing on patient assessment of pain and of risk for abuse, misuse or addiction; patient monitoring to evaluate treatment efficacy; and best practices for clinicians in the field of pain management

Calendar of Live Meetings and Events

A series of accredited regional meetings for pain management clinicians, focusing on balancing effective opioid therapy with appropriate assessment and monitoring for potential risk of abuse, misuse and addiction



Breakthrough News

A rotating series of headlines and daily news articles highlighting breaking news in the field of pain management

Clinical Expert Commentary

Leading experts provide their interpretation of hot topics in the field of pain management

Summaries of the latest clinical journal articles relevant to the field of pain management

A nonbranded educational website





ASAM CONFERENCE CALENDAR

ASAM -

May 4, 2006

Ruth Fox Course for Physicians San Diego Sheraton Hotel & Marina San Diego, California [8 Category 1 CME Credits]

May 5-7, 2006

37th Annual Medical-Scientific Conference San Diego Sheraton Hotel & Marina San Diego, California [21 Category 1 CME Credits]

July 21-23, 2006

Medical Review Officer (MRO) Training Course (Basic) Ritz Carlton Phoenix Hotel Phoenix, Arizona [8 Category 1 CME Credits]

October 26-28, 2006

ASAM Review Course in Addiction Medicine Westin O'Hare Hotel Chicago, Illinois [21 Category 1 CME Credits]

October 29, 2006

Course on Pain & Addiction Westin O'Hare Hotel Chicago, Illinois [8 Category 1 CME Credits]

December 8-10, 2006

Medical Review Officer (MRO) Training Course (Basic & Advanced) Marriott Metro Center Hotel Washington, DC [8 Category 1 CME Credits]

OTHER EVENTS OF NOTE

April 22-26, 2006

2006 AATOD National Conference: "Treating People With Dignity — Working With Criminal Justice and Health Care Systems" American Association for the Treatment of Opioid Dependence Hyatt Regency Atlanta, Georgia [Visit: www.AATOD.org or email AATOD@TALLEY.COM]

April 28-29, 2006

Second Annual National
Aging and Addiction Conference
Marriott West Palm Beach Hotel, West
Palm Beach, Florida
Sponsored by the Hanley Center
[Visit: www.hanleycenter.org/naac]

Except where otherwise indicated, additional information is available on the ASAM web site (www.ASAM.ORG) or from the ASAM Department of Meetings and Conferences at 4601 No. Park Ave., Suite 101, Chevy Chase, MD 20815-4520; phone 301/656-3920; fax 301/656-3815; email EMAIL@ASAM.ORG.

BUPRENORPHINE TRAINING -

For information or to register for a Buprenorphine Training Course, contact 1-888/362-6784 or visit the website: WWW.DOCOPTIN.COM.
All courses are approved for 8 Category 1 CME credits.

April 22, 2006

Atlanta, George Sponsored by ASAM & the Georgia Society of Addiction Medicine

May 6, 2006

Augusta, Maine Sponsored by ASAM & the Maine Society of Addiction Medicine

May 7, 2006

San Diego, California Sponsored by ASAM & the California Society of Addiction Medicine

May 10, 2006

Madison, Wisconsin
Sponsored by ASAM & the
Wisconsin Society of Addiction
Medicine

June 3, 2006

Columbia, Maryland Sponsored by ASAM & the Maryland Society of Addiction Medicine

September 7, 2006 San Francisco, CA

Sponsored by ASAM & the California Medical Association

COMPONENT SESSIONS

The following Component Sessions, organized by ASAM councils and committees, are scheduled for 8:00 to 10:00 p.m. Thursday, May 4th during ASAM's Med-Sci Conference (see the final program for locations).

- "Opioid Agonist Treatment 2005 Training, Clinical Practice and Policy Initiatives"
- "Why Not Private Practice? Practice Management: Coding and Billing Strategies"
- ★ "Public Policy in Addiction Medicine: Past, Present, and Future"
- ★ "Addiction Medicine and Primary Care"
- ★ "Chapters Council / SMSS Program"

BUPRENORPHINE AND OFFICE-BASED TREATMENT OF OPIOID DEPENDENCE

Sunday, May 7, 2006 • 8:00 am – 5:30 pm • San Diego Sheraton Hotel & Marina • San Diego, California

Course Director: David Fiellin, M.D., Yale University Medical School

This course is designed for physicians who have an interest in or experience with treating opioid-dependent patients, and who wish to qualify to use buprenorphine in office-based treatment of opioid dependence.

Topics to be addressed by an expert faculty include:

- Overview of opioid dependence and rationale for opioid agonist treatment
- Legislative changes allowing office-based treatment
- General pharmacology of the opioids
- Pharmacology, efficacy and safety of buprenorphine and buprenorphine/naloxone
- Clinical uses of buprenorphine and buprenorphine/naloxone, including induction, maintenance, and pharmacologic withdrawal
- Patient assessment and selection
- Office procedures and logistics
- Medical comorbidities in opioid-dependent patients
- Psychiatric comorbidities in opioid-dependent patients
- The role of psychosocial counseling in the treatment of opioid dependence
- Special treatment populations, including adolescents, pregnant women, and pain patients

The course is approved for up to 8 credit hours of Category 1 continuing education credit. (Only those who attend the full 8-hour program are eligible for a certificate of attendance.)

A separate registration fee is required for this course.

ATTENDANCE IS LIMITED, SO BE SURE TO REGISTER EARLY!

Visit ASAM's web site at WWW.ASAM.ORG, or register on-site (registration opens at 7:15 a.m.).

EXHIBITORS OFFER NEW PRODUCTS, SERVICES

Organizations/
Associations/Agencies
Al-Anon Family Groups Booth 408 Alcoholics Anonymous Booth 210
American Academy of
Addiction Psychiatry Booth 108
American Society of Addiction Medicine Ballroom Foyer
California Society of Addiction
Medicine Booth 110 Narcotics Anonymous
World Services Booth 305
National Alliance of Advocates for Buprenorphine Treatment Booth 116
National Institute on Alcohol Abuse and Alcoholism Booth 306
National Institute on Drug Abuse Booth 304
Ruth Fox Memorial Endowment Fund Display
Substance Abuse and Mental Health
Services Administration Booth 308
Book and Software Publishers
Blackwell Publishing Ltd Booth 212
Cold Spring Harbor Laboratory Press
Elsevier Services Booth 201
Lippincott Williams
& Wilkins Booth 602
S. Karger, AGDisplay
Pharmaceutical Manufacturers and Other Products & Services
Ameritox, Ltd Booth 300
Cephalon & Alkermes Booths 101-103
Clinical Tools, Inc Booth 506
D2 Market Research Booth 310
Forest Pharmaceuticals, Inc Booth 206
Mallinckrodt
Pharmaceuticals
NOVX Systems, Inc Booth 508 Odyssey
Pharmaceuticals, Inc Booth 603
OraSure Technologies, Inc Booth 302
Reckitt-Benckeiser Pharmaceuticals Booths 200-202
Sebia Electrophoresis Booth 404
United States Drug Testing
Laboratories Booth 102
VistaPharm, Inc Booth 511
Treatment Programs
Abundant Life and Adolescent Growth, Inc Booth 600
Addiction Recovery Resources
of Now Orloans Booth 410
of New Orleans Booth 410
Amity Circle Tree Ranch Booth 510
Amity Circle Tree Ranch Booth 510 Bayside Marin Booths 507-509
Amity Circle Tree Ranch Booth 510
Amity Circle Tree Ranch Booth 510 Bayside Marin Booths 507-509 Betty Ford Center at

rkuvulis, sek	VICES
Caron Foundation	Booth 113
Cedar — University of Colorado Hospital	Booth 606
Center for Professional	
Excellence Central Recovery	Booth 104
Treatment	Booth 207
COPAC, Inc	Booth 407
Cornerstone of Southern California	Booth 116
CRC Health Group	
Cri-Help, Inc.	
Crossroads Centre, Antigua	Booth 307
Eating Disorders Center	D+/- 200
of Denver Edgewood	
Elmhurst Memorial Healthc	
Professionals At Risk	
Treatment Emerging Solutions in Pain	
ROI Medical Group Boot	
Father Martin's Ashley	Booth 500
Friends Research Institute, Inc	Booth 301
Hazelden — Hazelden	
Springbrook	
Hythiam	
Lasting Recovery Outpatien Treatment Program	
Life's Journey Center	
Little Hill Foundation, Inc	Booth 413
Marworth Treatment Center	Booth 211
Memorial Hermann Prevent Recovery Center	
Michael's House — The Trea Center for Men	tment Booth 309
New York — Presbyterian Hospital	Booth 610
Palmetto Addiction	
Recovery Center Pine Grove — Next Step	
Practical Recovery Services	
Progressive Health Center.	
Promises Treatment Center	
Refuge, A Healing Place	Booth 114
Remuda Ranch	Booth 402
Ridgeview Institute, Inc	Booth 109
Rush Behavioral Health	Booth 313
Santé Center for Healing	
The Society for the Advance of Sexual Health	
Sequoia Center	Booth 400
Shepherd Hill	
Talbott Recovery Campus	Booth 412
University of Alabama at Birmingham — Addiction Recovery Program	Rooth 602
Valley Hope Association	
Williamsburg Place & the	
William J. Farley Center	DOULT 409

NIDA SYMPOSIUM FEATURES CLINICAL APPROACHES TO HIV AND HCV

FRIDAY, 9:30 a.m. – 12:30 p.m. and 2:30 p.m. – 6:00 p.m.

HARBOR ISLAND 2 & 3

Faculty: Andrea Cox, M.D., Ph.D.; Brian Edlin, M.D., M.P.H.; Richard Garfein, Ph.D., M.P.H.; Ramesh Ganju, Ph.D.; Kristy Marie Hendricks, Sc.D.; Charles B. Hinkin, Ph.D., ABPP; Robert Horsburgh, M.D.; Kenneth E. Sherman, M.D., Ph.D.; Jack Stapleton, M.D.; Ellen Tedaldi, M.D. and Terry Wright, M.D.

The National Institute on Drug Abuse (NIDA) has organized a day-long symposium, "Clinical Approaches to HIV and Hepatitis C Infections in Drug Abusers," to address issues important to addiction specialists who care for patients at risk for or diagnosed with these common infectious diseases.

An expert faculty will discuss the epidemiology and pathophysiology of HIV and HCV in drug-using populations, neuropsychiatric and psychological complications, and new medications for the treatment of the addicted and infected patient. Speakers also will discuss gaps in the knowledge base and outline directions for future research.

The symposium has been organized by Frank J. Vocci, Jr., Ph.D., and Jag H. Khalsa, Ph.D., of NIDA's Division of Medications Development.

NIAAA SYMPOSIUM HIGHLIGHTS **NEW MEDICATIONS FOR ALCOHOLISM**

SATURDAY, 10:00 a.m. - 12:00 noon and 2:30 p.m. - 5:30 p.m.

HARBOR ISLAND 2 & 3

Faculty: Raymond F. Anton, M.D.; Jack R. Cornelius, M.D., M.P.H.; Bankole A. Johnson, M.D., Ph.D.; George F. Koob, Ph.D.; Henry R. Kranzler, M.D.; John M. Littleton, M.D., Ph.D.; Barbara J. Mason, Ph.D. and Mark L. Willenbring, M.D.

A comprehensive overview of medications for the treatment of alcoholism is the focus of a symposium organized by the National Institute on Alcohol Abuse and Alcoholism (NIAAA). Faculty for "Medications Development for Alcoholism: From the Bench to the Patient," will offer a multidisciplinary perspective on medications development, highlighting the translation of drug discoveries through preclinical testing and clinical trials to adoption in clinical practice.

Organized by Mark L. Willenbring, M.D., and Raye Z. Litten, Ph.D., the symposium will present up-to-the-moment information on:

- Preclinical advances in discovering and testing novel compounds.
- The results of clinical trials of naltrexone, topiramate, ondansetron and other promising medications.
- New research findings on the treatment of adults and adolescents with psychiatric comorbidities.
- The role of genetics in predicting treatment outcomes with various medications.
- Real-world issues that confront physicians who would use medications in clinical practice.

STATE CHAPTERS TO MEET

In addition to the SMSS/Chapters Council meeting scheduled for Thursday evening, May 4th, from 7:00 to 10:00 p.m., the following ASAM Chapters and State Societies have scheduled meetings during Med-Sci. See the staff at the ASAM Registration Desk for meeting room locations.

Arizona	Saturday, May 6th, 9:00 a.m.
California	Saturday, May 6th, 4:30 p.m.
Colorado	Saturday, May 6th, 12:30 p.m.
Florida	Saturday, May 6th, 2:00 p.m.
Illinois	Friday, May 5th, 12:30 p.m.
Maryland	Friday, May 5th, 6:00 p.m.
Massachusetts	Friday, May 5th, 4:00 p.m.
Oregon	Friday, May 5th, 7:00 p.m.
New York	Friday, May 5th, 12:30 p.m.
Northwest	Friday, May 5th, 10:00 a.m.
Region VII	Friday, May 5th, 2:00 p.m.
Region IX	Friday, May 5th, 5:00 p.m.
Tennessee	Friday, May 5th, 12:30 p.m.
Texas	Friday, May 5th, 8:00 p.m.

MED-SCI FUNDERS ACKNOWLEDGED

ASAM is grateful to the following individuals and organizations for their generous support of this year's Medical-Scientific Conference:

Bendiner & Schlesinger **Bradford Health Services** California Society of Addiction Medicine **Center for Substance Abuse Treatment** Cephalon, Inc. COPAC, Inc.

Dr. and Mrs. Joseph E. Dorsey Endo Pharmaceuticals, Inc. The William J. Farley Center Forest Pharmaceuticals, Inc. Hazelden & Hazelden — Springbrook John P. McGovern Foundation Marworth Treatment Center **Memorial Hermann Prevention** & Recovery Center Pine Grove/Next Step Ridgeview Institute Santé Center for Healing Schick Shadel Hospital **Christopher D. Smithers Foundation**

Tyco Healthcare — Mallinckrodt, Inc.

CSAT SYMPOSIUM TO FOCUS ON METHAMPHETAMINE

SUNDAY, May 7, 2006 * 8:30 a.m. to 12:30 p.m.

GRAND BALLROOM B



Faculty: H. Westley Clark, M.D., J.D., M.P.H., FASAM; Rachel Gonzales, Ph.D.; William Haning, M.D.; Walter Ling, M.D.; Edyth London, Ph.D.; Jane Maxwell, Ph.D.; Richard Rawson, Ph.D.; Shirley Semple, Ph.D.; Steve Shoptaw, Ph.D. and Matt Torrington, M.D.

To provide ASAM members with the latest information on methamphetamine addiction and its treatment, the Center for Substance Abuse Treatment (CSAT) will bring together some of the nation's leading experts on the drug for a half-day symposium on Sunday, May 7th.

Chaired by CSAT Director H. Westley Clark, M.D., J.D., M.P.H., FASAM, "Treatment of Methamphetamine Use Disorders" was organized by Anton C. Bizzell, M.D., Medical Director in CSAT's Division of Pharmacologic Therapies. Faculty members will address the following topics:

- · Epidemiology of methamphetamine abuse;
- Methamphetamine's effects on the brain and behavior;
- Medical aspects of methamphetamine use disorders;
- Methamphetamine withdrawal and its management;
- Identification and management of psychological complications;
- Psychosocial and behavioral treatments: The Matrix Model;
- Treatment issues for special groups, including women and adolescents.

